

Symmetry Breaking: A New Paradigm for Non-Perturbative QFT and Topological Transitions

© 2009 by [Daniel D. Ferrante](#)

[Physics Department, Brown University](#)
Providence — RI. 02912. USA.

Preface

Lately, after ~ 50 years, there seems to be a convergence in the languages used to describe Quantum Field Theory and String Theory, such that it seems possible to relate objects from these two perspectives. Therefore, a deeper investigation of the properties and features of QFT is a reasonable thing to do: non-perturbative effects, dualities, emergent properties, non-commutative structures, etc. This particular line of research uses Symmetry Breaking in order to probe a few of the different topics mentioned above, i.e., Symmetry Breaking is used as an “underlying principle”, bringing different features of QFT to the foreground. However, the understanding of Symmetry Breaking that is used here is quite different from what is done in the mainstream: Symmetry Breaking is understood as the solution set of a given QFT, its vacuum manifold, or, more modernly, its Moduli Space. Distinct solutions correspond to different sectors, phases, of the theory, which are nothing but distinct foliations of the vacuum manifold, or points in the Moduli Space (for all possible values of the parameters of the theory). Under this framework, three different problems will be attacked: “Mollifying QFT”, “Topological Transitions and Geometric Langlands Duality” and “Three-dimensional Gravity and its Phase Transitions”. The first makes use of the Moduli Space of the theory in order to construct an appropriate mollification of it, rendering it viable to simulate a QFT in Lorentzian spaces, tackling the “sign problem” heads-on. The connections with Lee-Yang zeros and Stokes Phenomena will be made clear. The second will show that each different phase has its own topology which can be used as Superselection Rule; moreover, the Euler Characteristic of each phase gives it quantization condition. The mechanism via which several dualities work will also be elucidated. The last one will generalize a 0-dimensional QFT, via dimensional construction through its D -Module, and conjecture several connections between the Lie-algebra-valued extension of the Airy function and the recent Partition Function found for three-dimensional gravity with a negative cosmological constant. These three problems, put together, should exhibit a solid and robust framework for treating QFT under this new paradigm.

Acknowledgements

The author would like to thank his advisor, [G. Guralnik](#), for invaluable support, Physics- and other-wise.

Further thanks go to Professor [Michael Kosterlitz](#) and [James O'Dell](#), for discussions about Physics (including phase transitions, non-equilibrium statistical mechanics, symmetry breaking and renormalization methods), Scientific and High Performance Computing.

Also, the author would like to thank Professors [Herbert Fried](#), [Bruno Harris](#), [John Savage](#), [James Baird](#), [Ian Dell'Antonio](#), [Antal Jevicki](#), and [Marcus Spradlin](#), for all forms of discussions (academic or not) and various “coffee and cookies” moments.

Finally, [C. Pehlevan](#) is acknowledged for discussions and useful conversations. This work was supported in part by funds provided by the US Department of Energy (DOE) under contracts DE-FG02-85ER40237 and DE-FG02-91ER40688-TaskD.

Contents

1. Motivations and Introductory Remarks	1
1.1. The road ahead	4
I. Mollifying Quantum Field Theory	7
2. Motivation and Introduction	9
2.1. Sign Problem	9
2.2. Smoothing out the Measure	10
3. Mollifying Quantum Field Theory	11
3.1. Importance Sampling	12
3.2. Mollifying the Importance Sampling Function	13
4. Tuning the Mollification	17
5. Simulation Details	19
5.1. Random Number Generators	19
5.2. Non-local Algorithms	19
5.3. Euclidean Lattice Simulations	20
5.4. Mollified Minkowski Lattice Simulations	21
6. Spontaneous Symmetry Breaking	23
7. Results	27
7.1. Lower Dimensional	27
7.1.1. Airy Function	27
7.1.2. 0-dimensional ϕ^4 Theory	31
7.2. Higher Dimensional	35
II. Solution Space and Moduli Space Topology	37

8. Introduction	39
9. Classical Field Theory and the Jacobi Metric	41
9.1. Harmonic Oscillator	42
9.2. Cubic Potential	44
10. Applications in Quantum Field Theory	47
10.1. The $\lambda \phi^4$ Potential	51
11. Gauge Theory Examples	55
11.1. The Landau-Ginzburg Functional	56
11.2. The Seiberg-Witten Functional	59
12. Conclusions	63
 III. Three-Dimensional Gravity and Airy Functions	 65
13. Introduction	67
14. Extended Airy Functions	71
15. Dimensional Construction Sketch	73
 IV. Final Conclusions	 75
16. Conclusions and Future Work	77
 V. Appendices	 81
A. Convolutions and Smoothing	83
A.1. Mollifiers	83
A.2. Properties of Mollifiers	84
B. Entropy Calculations	85
C. Parabolic Cylinder Functions	87
 VI. Bibliography	 89

Motivations and Introductory Remarks

The paradigm of research in Physics and Mathematics around the 1960's was that Physics dealt with local structures (QFT, “Dual Models”, etc) while Mathematics dealt with global tools (Morse Theory, Cohomologies, Index Theorems, etc).

Nowadays, this situation is somewhat reversed, with physicists worrying about questions of Yang-Mills over Riemann surfaces, Higgs Bundles, Topological Quantization, etc; while mathematicians have gotten their hands full with structures inspired by QFT and String Theory, e.g., D -Modules.

Thus, it seems to be that there is a convergence of these pictures into a single language: after ~ 50 years it looks possible to translate between these two pictures, i.e., it seems possible to relate objects from these two perspectives.

Therefore, a deeper investigation into the properties and features of QFT seems a reasonable thing to do. Studies in dualities (AdS/CFT, Langlands), emergent properties (effective theories), non-commutative structures, non-perturbative effects (branes), etc, seem like a rich venue to follow.

The present work started as a means to deepen the understanding between symmetry breaking and the multiple solutions¹ of a certain quantum field theory, in the spirit already developed in [35].

To better understand this line of argument, let us focus on 0-dimensional (i.e. ultralocal) models for the sake of simplicity. The idea is to compute the Schwinger-Dyson Equations (SDEs) of the theory in question and solve it analytically. Generically speaking, given that most theories are defined by a Potential which is a polynomial in the fields, the SDEs are expected to have more than one solution. Therefore, not all of these solutions will admit a series expansion in terms of the coupling constants (which are normally called “perturbative series”), implying that some solutions are non-perturbative, even though they admit a series expansion in some other parameter. In this sense, some of these solutions may keep the original symmetries (present in the Action), but some may not. To fix the idea, let us look at two particular examples.

¹Henceforth, when referring to QFT and its solutions, we shall use interchangeably the following terms: *solution space*, *configuration/phase space*, *vacuum manifold* and *moduli space*.

Airy Potential This theory is defined (in 0-dimensions) by $S(\phi) = \phi^3/3 + J\phi$, and its Feynman Path Integral (FPI) representation is given by,

$$\mathcal{Z}(J) = \frac{\int_{\mathbb{C}} e^{i\frac{\phi^3}{3} + iJ\phi} d\phi}{\int_{\mathbb{C}} e^{i\frac{\phi^3}{3}} d\phi} < \infty ;$$

where $\mathbb{C}$ is the contour of integration that renders this FPI finite. On the other hand, its SDE is obtained in the following way:

$$(\phi^2 - J) \mathcal{Z}(J) = 0 \xrightarrow{\phi \mapsto -i\partial/\partial J} (\partial_J^2 - J) \mathcal{Z}(J) = 0 .$$

Now we make an important observation: the SDE above is second order and, thus, must have two solutions. As it turns out, these solutions are given by the $\text{Ai}(J)$ and $\text{Bi}(J)$ functions. At this point, it is not difficult to realize that the FPI is nothing but the integral representation of the differential problem posed above, which implies that,

$$\mathcal{Z}[J] = \frac{\int_{\mathbb{C}_{1,2}} e^{i\phi^3/3 + iJ\phi} d\phi}{\int_{\mathbb{C}_{1,2}} e^{i\phi^3/3} d\phi} = \text{Ai}[J] \text{ or } \text{Bi}[J] ;$$

where the contours $\mathbb{C}_{1,2}$ are such that the integral representation above is finite.

The realization that should be accomplished at this stage is that ϕ -space had to be “complexified” so that we would be able to find both of these solutions — usually, when we think in terms of a FPI, it is *presumed* that the integration above should be done for $\phi \in \mathbb{R}$. However, as we just showed above, in this very simple example we already see that if we do not extend the FPI representation to $\phi \in \mathbb{C}$, we will not obtain *all* of the possible solutions to the problem at hand.

Having said that, it is important to realize that the partition function, in its most general form, will be given by a linear combination of all possible solutions, i.e.,

$$\mathcal{Z}(J) = \alpha \text{Ai}(J) + \beta \text{Bi}(J) ;$$

where $\alpha, \beta \in \mathbb{C}$ are scalars.

Further, the asymptotic series of $\text{Ai}(J)$ and $\text{Bi}(J)$ has a different form in different quadrants of the complex plane, a fact known as the Stokes phenomenon, implying that $\mathcal{Z}(J)$ will be a meromorphic function. Therefore, resummations of these asymptotic series, if it has to be done at all, must be tackled with the utmost care and attention, once mixing different branches of these series renders all results obtained after that completely useless.

Also, because $J \in \mathbb{C}$, we can study the partition function in terms of its properties with respect to the Möbius group, which is the automorphism group of the Riemann sphere. In particular, we may focus in one of its more important subgroups, the Modular Group ($\text{SL}(2, \mathbb{Z})$, where we identify the elements A and $-A$), and study all of the connections with modular forms, elliptic curves and many fractals — some

connections with String Theory can be drawn once it is realized that the generators of this group are given by “unit translation to the right” and “inversion in the unit circle followed by reflection about the line $\Re(z) = 0$ ”, where this last generator is analogous to T-duality (which can be understood in terms of Mirror Symmetry).

This is an interesting example because the cubic potential is usually said to be unbounded from below and, therefore, yields no meaningful theory. However, we clearly see above two solutions which are *finite*. This happens because we are explicitly looking for boundary conditions to solve the SDE, a fact that is completely analogous to searching for contours that render the Path Integral finite.

Furthermore, this example can be readily extended from a scalar-valued field ϕ , to more interesting cases such as that of a matrix-valued field Φ , or of a Lie-algebra valued field φ . As expected, the only modification to the above equations is that the Airy function is extended analogously to either a matrix-valued or Lie-algebra-valued function — in fact, these extensions are *defined* via the FPI representation above [53].

Once these extensions are established, going from 0-dim to d -dim is accomplished via the use of D -Modules: in this sense, the D -Modules are used to compute the particular “Thermodynamic Limit” arising from the 0-dim theory at hand. This will be important for future developments.

It is important to emphasize that the *observables* of a theory should be real, once they are the measurable quantities. But this says nothing about the nature of the partition function nor of the fields themselves.

$\lambda \phi^4$ Potential This theory is defined by $S(\phi) = \mu \phi^2/2 + \lambda \phi^4/4$. The SDE and FPI for this model are:

$$(\mu \phi + \lambda \phi^3 - J) \mathcal{Z}[J] = 0 \xrightarrow[g=\mu/\lambda]{\phi \mapsto -i \partial/\partial J} (\partial_J^3 + g \partial_J - J) \mathcal{Z}[J] = 0 ;$$

$$\mathcal{Z}[J] = \frac{\int_{\mathbb{C}_{1,2,3}} e^{i(\mu \phi^2/2 + \lambda \phi^4/4 - J \phi)} d\phi}{\int_{\mathbb{C}_{1,2,3}} e^{i(\mu \phi^2/2 + \lambda \phi^4/4)} d\phi} = U[g, J] \text{ or } V[g, J] \text{ or } W[g, J] ;$$

where $g = \mu/\lambda$, and the three solutions for the SDE are given by the Parabolic Cylinder Functions, $U[g, J]$, $V[g, J]$ and $W[g, J]$, which, in turn, can be obtained with three different contours, $\mathbb{C}_{1,2,3}$, in the FPI representation (see Appendix C).

As mentioned earlier, each one of these parabolic cylinder functions represents one particular solution: symmetric, broken-symmetric and solitonic.

Again, all of the considerations made above can be readily applied in this case: complexification of ϕ -space in order to obtain all possible solutions, the partition function is a linear combination of these three solutions, Stokes phenomenon is present in the asymptotic series for each one of these solutions, the partition function is meromorphic and, finally, its properties with respect to the Modular Group can be

studied.

1.1. The road ahead

We have just shown how extremely important boundary conditions (resp. contours) are for properly defining a QFT and all of its solutions/sectors (resp. moduli space); and this is the cornerstone of our work: from this point on, we can attack several different problems in novel ways.

Here are the problems that we tackle in this thesis:

Mollifying QFT We probe these multiple solutions using the help of a mollifier, which is essentially a band-pass filter that eliminates the meaningless highly-oscillatory parts of the Partition Function, but does retain its meaningful structure. This shows clearly the links with Lee-Yang Zeros (complex zeros of the Partition Function, seen as a polynomial in the coupling constants) and Stokes phenomena (the asymptotic expansion of each solution has a parameter-dependent limit, which means that there is a discontinuous change [in the Partition Function] when crossing Stokes lines separating distinct solutions/sectors of the theory). For more on this, please see [40].

Topological Transitions and Geometric Langlands Duality We show that each different solution/sector of the theory has a distinct topology which can be used as a Superselection Rule (topological charge); moreover, the Euler Characteristic (second Chern class) of each phase gives its quantization condition. Also, these sectors can be labeled by the discriminant of a polynomial constructed from the potential, where the structure of this discriminant is related to singularities in the Partition Function, which, in turn, must be treated as a meromorphic function (which connects with the picture above about Lee-Yang zeros and Stokes Phenomena). However, the different solutions of the same theory are related by “dualities” in the values of the parameters of the theory: distinct boundary conditions establish different allowed ranges for these parameters. In this sense it is possible to construct several “dualities” that translate one sector into another, and this is how we make touch with the subject of Geometric Langlands Duality. Furthermore, using D -Modules, we can take the Thermodynamic Limit of each particular 0-dim theory into its full d -dim version, thus dimensionally constructing each different sector of the theory: each branch is “grown” out of the “natural differential operator” defined by the specific potential and its parameters (determined by the appropriate boundary conditions). Therefore, these dualities are not only with respect to different sectors of the theory, but also with respect to each natural differential operator associated to each phase, thus establishing the possible allowed values for the geometric quantities (connection) associated to each point in the moduli space of the whole theory. For more on this, please refer to [41, 55].

Three-dim Gravity and Phase Transitions This is where we make use of the extension to Lie-algebra-valued fields mentioned above when solving the Airy potential.

Recently, Witten has shown that 3-dim gravity with negative cosmological constant has a dual 2-dim CFT, [39]. In order to do so and obtain a holomorphically factorized Partition Function, he had to move from real saddle points (which do not account for the whole theory) to *complex* ones. And, in doing so, he found that the Hawking-Page phase transition is analogous to the condensation of Lee-Yang zeros. The question we are tackling is that of extending the Airy function as previously discussed (to a Lie-algebra valued Airy function) and “grow” this 0-dim solution into a 3-dim one using its D -Module. Furthermore, there are some questions that arise from the connection with Braid and Knot theory, such as whether having two distinct Partition Functions (for A_i and B_i) imply different polynomial invariants, or whether this implies that the structure constants of the Lie group in question can be dynamically determined (by the allowed contours), etc.

Part I.

Mollifying Quantum Field Theory

Motivation and Introduction

2.1. Sign Problem

One of the fundamental difficulties of Monte Carlo (MC) approaches is known as the “sign problem”. It is encountered when the functional integrals to be evaluated do not have a positive definite measure. It is not related to any approximations or fundamental errors in the MC scheme, but it describes the situation where the statistical error can become very large. In general, any expectation value can be written as,

$$\langle O \rangle = \frac{\int O[\phi] \mu[\phi]}{\int \mu[\phi]}, \quad (2.1)$$

where μ and O are real-valued functions of the field variables and, in general, the measure $\mu[\phi]$ need not be positive: in Lorentzian QFT this measure is complex-valued and given by $e^{iS[\phi]} \mathcal{D}\phi$. However, if $\mu[\phi]$ changes sign, it cannot be considered a probability density (*i.e.*, a measure).

The standard trick to avoid this problem is to modify the measure in the following way: $\tilde{\mu}[\phi] = |\mu[\phi]| / \int |\mu[\phi]|$. Then, one absorbs the sign of $\mu[\phi]$ in the quantity to be measured:

$$\langle O \rangle = \frac{\int O[\phi] \text{sign}(\mu[\phi]) \tilde{\mu}[\phi]}{\int \text{sign}(\mu[\phi]) \tilde{\mu}[\phi]}. \quad (2.2)$$

In some cases this may work. However, the random walk guided by $|\mu[\phi]|$ is very likely to predominantly sample unimportant regions in phase space.

At this time no completely satisfactory solution to the sign problem exists, although there are some promising attempts including positive projection [2, 3], fractal decomposition scheme [4], and Berry’s phase [5] and its Stiefel manifold [6]. We believe that the techniques discussed in the present work lay the grounds for another possible solution path. The general success of the methods studied here will depend

on the development of more computationally efficient algorithms than the ones used to illustrate the simple examples treated below.

2.2. Smoothing out the Measure

The idea behind the mollification technique is to use a convolution in order to smooth out and filter the measure: the highly oscillatory measure given by $e^{iS[\phi]} \mathcal{D}\phi$ is convoluted with some suitable function (called a *mollifier*; see Appendix A for more details) and as a result only an effective contribution is left (rapid oscillations of the measure cancel out when integrated over the slowly varying mollifier).

A convolution is an integral that expresses the amount of overlap of one function g as it is shifted over another function f . It therefore “blends” one function with another. In mathematics, mollifiers are smooth functions with special properties, used in distribution theory (generalized functions) to create a sequence of smooth functions approximating non-smooth functions, via a convolution.

In an application of this idea to QFT we will replace the rapidly oscillating measure in Minkowski space ($I = \exp\{iS[\phi]\}$) by smooth functions ($I_\epsilon = \eta_\epsilon * I$) so that we can run numerical simulations that are otherwise prohibitive (see A). At this stage, the only requirement that is made is that: $\int \eta_\epsilon(x) dx = 1$, where the integral is performed over the domain of η_ϵ . The parameter ϵ controls the approximation, the smoothness, being made, and in the $\epsilon \rightarrow 0$ limit, the original expressions are recovered. (The meaning of this ϵ parameter is that of dilating or contracting the mollifier itself, thus controlling the range over which the filtering is done. Note that the calculations should be done using a non-vanishing value for ϵ in order to avoid the analytical expressions which are hard to handle numerically.)

Mollifying Quantum Field Theory

The technique described above (and detailed in [A](#)) can be applied to [Lattice] QFT in the following way:

$$\mathcal{Z}_\epsilon[J] \equiv \int \left\{ \int \eta_\epsilon[\phi - \varphi] e^{iS[\varphi;J]} \mathcal{D}\varphi \right\} \mathcal{D}\phi . \quad (3.1)$$

This form is useful to set up numerical computations. Performing the mollification before performing the path integral can be very advantageous.

Mollifying the integrand (i.e., the complex exponential of the action) smooths the highly oscillatory integral. The integrand changes its form, from the canonical I to the mollified I_ϵ , in the following fashion,

$$I[\phi; J] = e^{iS[\phi;J]} \longmapsto I_\epsilon[\varphi; J] = (\eta_\epsilon * I)[\varphi; J] = \int \eta_\epsilon[\varphi - \phi] e^{iS[\phi;J]} \mathcal{D}\phi ,$$

i.e., the convolution with η_ϵ changes variables: $\phi \mapsto \varphi$. Taking [functional] derivatives of $\mathcal{Z}$ (which yield Green's functions) is just the same as taking them with respect to $\mathcal{Z}_\epsilon$, because the derivative operator commutes with the mollification.

The [Feynman] Path Integral is constructed just as before:

$$\begin{aligned} \mathcal{Z}[J] &= \int I[\phi; J] \mathcal{D}\phi ; \\ \mathcal{Z}_\epsilon[J] &= \int I_\epsilon[\varphi; J] \mathcal{D}\varphi . \end{aligned}$$

Note that,

$$\begin{aligned} \mathcal{Z}_\epsilon[J] &= \int \left\{ \underbrace{\int \eta_\epsilon[\varphi - \phi] e^{iS[\phi;J]} \mathcal{D}\phi}_{I_\epsilon[\varphi;J]} \right\} \mathcal{D}\varphi . \\ \therefore \mathcal{Z}_\epsilon[J] &\equiv \mathcal{Z}[J] ; \end{aligned} \quad (3.2)$$

where standard properties of convolutions (see [7]) have been used in the last step.

Even though the above result is analytic, when one goes to the simulations, a small dependence on ϵ shows up. (See Section 4 for a more detailed discussion on this matter.)

3.1. Importance Sampling

The next step consists of choosing an appropriate sampling function (for the Monte Carlo simulation):

$$\mathcal{Z}_\epsilon[J] \equiv \int \left\{ \int \eta_\epsilon[\phi - \varphi] \frac{e^{iS[\varphi; J]}}{W[\varphi]} \mathcal{D}W \right\} \mathcal{D}\phi ,$$

where $\mathcal{D}W = W[\varphi] \mathcal{D}\varphi$, i.e., just a change of variables in order to make the computations more robust: it is possible to use the integrand profile in order to speed up the calculation.

An appropriate choice for the importance sampling function, W , is usually given by $W_\epsilon[\phi] = |I_\epsilon[\phi]|$, [8].

A little digression is in order: If all we wanted to do was to simulate a certain [Euclidean] QFT on a lattice, the above reasoning would be just as valid, modulo the mollification, i.e., we would conclude that the proper importance sampling function was $W[\phi] = |I[\phi]|$. The full-fledged formula is useless for doing computations since it implies full knowledge of the theory being calculated. A better choice is a simple approximation: we choose a saddle-point approximation in order to make the expression more manageable and still keep most of the characteristics of the integrand.

Thus, the importance sampling function ($W[\phi]$), which is used just to better guide the simulation, would be nothing but the absolute value of the sum over all saddle-points of the theory in question. Assuming we have only one such saddle-point, the well known answer is given by:

$$\begin{aligned} W[\phi] &\approx \left| \frac{e^{iS[\phi_0]}}{\sqrt{\det\{-\partial^2 + m^2 + \mathcal{V}''[\phi_0]\}}} \right| ; \\ &= \left| e^{iS[\phi_0]} \right| \left| e^{-\frac{1}{2} \text{tr}\{\log(-\partial^2 + m^2 + \mathcal{V}''[\phi_0])\}} \right| ; \\ &= e^{-\frac{1}{2} \text{tr}\{\log(-\partial^2 + m^2 + \mathcal{V}''[\phi_0])\}} \equiv \left(\det\{-\partial^2 + m^2 + \mathcal{V}''[\phi_0]\} \right)^{-1/2} . \end{aligned} \quad (3.3)$$

Using this to find an importance sampling function for a simple scalar QFT on the lattice requires solving for the determinant above. In Lattice QCD, the problem of the *fermion determinant* is a very time-consuming operation that must be done for *every* step of the MC calculation.

But this is not the only computational bottleneck. The more severe one comes from the so called *sign problem*: The exponent in $I[\phi]$ is not bounded from below, therefore one cannot guarantee the ergodicity of the Markov Chain underlying the

Monte Carlo draws, i.e., a much bigger number of draws would be needed in order to yield any meaningful answer. This is the reason why calculations are done in Euclidean space rather than Lorentzian/Minkowski space. In Euclidean space (Wick-rotating the integrand), the exponent in question becomes $I_E[\phi] = e^{-S[\phi]}$: it is bounded from below and the Monte Carlo method works fine (the Markov Chain behind it becomes ergodic).

It is to tackle this problem that the method of mollifiers comes in: mollifying $I[\phi]$ will yield a smooth function, $I_\epsilon[\phi]$, whose properties help the convergence of the MC computation. Furthermore, if this can really be accomplished and calculations with imaginary exponents become a reality, the next logical step is to analyze the different phases of the given QFT. As shown in [9], the different phases of a theory can be picked out with a mere choice of boundary conditions, which is the same as properly defining the measure of the [Feynman] Path Integral. The upshot is that if we want to obtain answers other than the ones that can be reached via perturbation theory, either the measure of the path integral or the boundary conditions of the Schwinger-Dyson equations have to be modified [9].

Once there is nothing preventing the measure of the [Feynman] Path Integral, from being (in the most general case) complex-valued — in order to account for the phase structure of the theory —, the fact that its integrand is also complex-valued has to be taken more seriously, afterall this is a non-perturbative result: a perturbative series only works if we know, *a priori*, in which phase we are working, so it can be tailored to that particular sector of the theory.

Thus, the need to address the sign problem becomes even clearer: it is not simply a problem of doing MC in Lorentzian/Minkowski space, it is mostly a problem of being able to compute all possible solutions — all the different phases — of a given QFT.

3.2. Mollifying the Importance Sampling Function

As outlined before, the idea is to use the mollification technique to handle the highly oscillatory terms present in the importance sampling function.

There are *three* possible choices for the sampling function W :

1. Taylor-expand and mollify the complex integrand: *no* knowledge about the saddle-points is necessary, but it is very computer intensive, which is the reason we will not focus on it;
2. As done in (3.5), where the integrand is saddle-point expanded and mollified, but the *explicit* form of the mollifier is used in order to perform the remaining integral: not so heavy on the computer, but some knowledge about the saddle-points is needed;
3. As done in (3.6), where the integrand is saddle-point expanded and mollified, generalizing the standard saddle-point approximation: less intensive of all 3, but even more knowledge about the saddle-points is needed.

The difference between the second and third methods above is that the integral in (3.4) is explicitly carried out in the former, but not in the latter. As explained below, the third method (listed above) consists of the simple generalization of the well known saddle-point approximation; however, we need to note that more knowledge about the particular saddle-points is needed in order to perform the contour integration involved in this scheme (and this is the difference with respect to the second method).

The calculation below shows the second method outlined above, known as $\mathcal{Q}^{nd}$ -order gradient approximation:

$$W_\epsilon[\phi] = |I_\epsilon[\phi]| = \left| \int \eta_\epsilon[\phi - \varphi] e^{i S[\varphi]} \mathcal{D}\varphi \right|.$$

Now, a saddle-point expansion is performed on the action,

$$W_\epsilon[\phi] \approx \left| \int \eta_\epsilon[\phi - \varphi] e^{i (S[\varphi_0] + \frac{1}{2} (\varphi - \varphi_0)^2 S''[\varphi_0])} \mathcal{D}\varphi \right|.$$

A mollifier needs to be chosen, and for the purposes of this calculation, a *Gaussian* one is our choice:

$$\eta_\epsilon[\phi - \varphi] = e^{-\frac{1}{2} (\phi - \varphi)^2 / \epsilon^2};$$

$$W_\epsilon[\phi] = \left| \int_{-\infty}^{\infty} e^{-\frac{1}{2} (\phi - \varphi)^2 / \epsilon^2} e^{i (S[\varphi_0] + \frac{1}{2} (\varphi - \varphi_0)^2 S''[\varphi_0])} \mathcal{D}\varphi \right|; \quad (3.4)$$

$$\Rightarrow W_\epsilon[\phi] = \left| \sqrt{2\pi\epsilon^2} \frac{\exp \left\{ i S[\varphi_0] + \frac{i}{2} \frac{(\phi - \varphi_0) S''[\varphi_0] (\phi - \varphi_0)}{1 - i \epsilon S''[\varphi_0] \epsilon} \right\}}{\sqrt{\det \{ 1 - i \epsilon S''[\varphi_0] \epsilon \}}} \right|. \quad (3.5)$$

The important point to note about the derivation above is that the *explicit* form of the mollifier, η_ϵ , had to be used and, because of that and the nature of the MC simulation, we do not necessarily need a good knowledge about φ_0 . This means that we can compute the above importance sampling function, (3.5), and use a *trial* φ_0^{trial} : the [MC] simulation will do the job of moving towards the exact φ_0 and pick out the different phases of the theory. As shown elsewhere, [11, 12, 13, 8], this works quite fine.

However, for more relevant cases (e.g. 4-dimensional QFTs) the time required for simulations becomes unrealistic. The bottom-line is that a simple-minded implementation of the method will have the code/computer doing the work of mapping the (ϕ, φ) -space, while one could use that information beforehand in order to speed things.

The way out of this is to generalize the following well known result: $\mathcal{J}(s) = \int_{\mathbb{C}} g(z) e^{s f(z)} dz = \sqrt{2\pi} g(z_0) e^{s f(z_0)} e^{i\alpha} / \sqrt{|s f''(z_0)|}$; $s \in \mathbb{R}$, $z \in \mathbb{C}$; i.e.,

$$W_\epsilon[\phi] = \left| \sqrt{2\pi} \frac{\eta_\epsilon[\phi - \varphi_0] e^{i S[\varphi_0]}}{\sqrt{|\det \{ S''[\varphi_0] \}|}} \right|. \quad (3.6)$$

At this point, the similarity between (3.3) and (3.6) is clear, in fact, they are the same formula, except that, in the former case there is no mollification while in the latter there is mollification.

Tuning the Mollification

At this point we are almost done with the analysis of the mollification technique, the only aspect remaining being the optimal choice of ϵ , the parameter that regulates the approximation.

From the discussion in appendix A, it is clear that in the $\epsilon \rightarrow 0$ limit we recover the original theory. However, numerically, this is not exactly what happens, because of numerical and statistical fluctuations. Therefore, the choice of an optimal value for ϵ is crucial.

Using an information theoretic viewpoint, we can choose ϵ to optimally compress $W_\epsilon[\phi]$ around each stationary phase point. Therefore, we can define an information entropy based on $W_\epsilon[\phi]$ in the following way:

$$\mathcal{S}[\epsilon] = - \frac{\int W_\epsilon[\phi] \log(W_\epsilon[\phi]) [d\phi]}{\left(\int W_\epsilon[\phi] [d\phi] \right) + \log \left(\int W_\epsilon[\phi] [d\phi] \right)}. \quad (4.1)$$

The optimal ϵ is the one that minimizes $\mathcal{S}[\epsilon]$. This value corresponds to maximal compression of the information in the importance sampling function.

In order to illustrate the above, let us consider a very simple [0-dimensional] example, given by $S[\phi] = m^2 \phi^2/2$. In this case, the saddle-point is $\phi_0 = 0$ and the importance sampling function and the information theoretic entropy are given by:

$$\begin{aligned} W_\epsilon[\phi] &= \frac{e^{-\phi^2/2\epsilon^2}}{\sqrt{2\pi\epsilon^2}} \frac{e^{iS[\phi_0]}}{\sqrt{S''[\phi_0]}}; \\ &= \frac{e^{-\phi^2/2\epsilon^2}}{m\epsilon\sqrt{2\pi}}; \\ \mathcal{S}[\epsilon] &= \frac{\log(m\epsilon\sqrt{2\pi}) - \epsilon^2}{1 - m\log(m)}. \end{aligned}$$

Plotted below are the graphs of the importance sampling function (also showing how it varies with ϵ) and of the information theoretic entropy (both using $m = 1$):




Figure 1: Importance Sampling function (leftmost plot, $m = 1$, $W_\epsilon[\phi] = e^{-\phi^2/2\epsilon^2}/\epsilon\sqrt{2\pi}$) and information theoretic entropy (rightmost plot, $m = 1$, $\mathcal{S}[\epsilon] = \log(\epsilon\sqrt{2\pi}) - \epsilon^2$) for the action $S[\phi] = m^2 \phi^2/2$.

We can clearly see the importance function peaking around $\phi = 0$, which is the expected behavior (*i.e.*, it happens around the saddle-point $\phi_0 = 0$), while the information theoretic entropy shows that the best parameter for the mollification is $\epsilon \rightarrow 0$: this is not desirable, once $\lim_{\epsilon \rightarrow 0} \mathcal{S}(\epsilon) \rightarrow -\infty$, but it simply shows that we should use a small non-vanishing value for ϵ , as mentioned before in Section 2.

Simulation Details

In order to perform the simulation as outlined in 5.4, some technical details had to be taken into more serious account. Among them, the important ones are: random-number generation and non-local algorithms. Those will be the bottle-necks, as noted in [14].

5.1. Random Number Generators

Even though some consider the random-number question a solved one, this is not always the case [14]. For this reason, the choice of which random-number generator (RNG) to use is still a critical one, not only for speed reasons but because of systematic errors as well. In order to address this issue the choice made was for the Mersenne Twister RNG [15] (period of $2^{19937} - 1$).

5.2. Non-local Algorithms

As for non-local algorithms, the choice made was for a Genetic Algorithm (GA) [16]. The “population”, in the Lattice QFT sense, consists of the lattice (*i.e.* spacetime) points. For a given “recombination rate”, an initial population gets *randomly* divided into pairs of [lattice] points and “crossing-over” (or “genetic”) operators are applied to these pairs in order to generate different sets of pairs. In this context, the crossing-over operators exchange the coordinates of given points, thus there are 4 of those operators, namely: T_t, T_x, T_y, T_z . That is, if the crossing-over operator T_x is applied to the points $\phi = (\phi_t, \phi_x, \phi_y, \phi_z)$ and $\phi' = (\phi'_t, \phi'_x, \phi'_y, \phi'_z)$ the outcome will be: $T_x[\phi, \phi'] = \{(\phi_t, \phi'_x, \phi_y, \phi_z), (\phi'_t, \phi_x, \phi'_y, \phi'_z)\}$. Analogous definitions are valid for the other crossing-over operators. The key features for this choice were: which T_i to use is random; whether or not the change $\phi \mapsto \phi'$ is made depends on a given probability and the recombination rate can be arbitrarily chosen (although keeping it below 1% showed to be a good tune). Moreover, the genetic operators are unitary, *i.e.*, $T_i = T_i^{-1}$. This implies that $T_i^2 = \mathbb{1}$, which guarantees the so-called *detailed balance* of the

MC simulation (this is the *ergodicity* of the algorithm). At this point, the only step remaining is the explanation of the *probability profile* used in algorithm 2 below.

- 1: Choose the recombination rate such that: $0 \leq r \leq 1$ {usually between 0.5% and 1.0%}
- 2: Draw a random number $\beta \in [0, 1]$ and compare with r
- 3: **if** $\beta \geq r$ **then**
- 4: Metropolis Monte Carlo loop {using the *random walk* technique}
- 5: **else**
- 6: Genetic loop {see algorithm 2 below}
- 7: **end if**

Algorithm 1: Genetic Algorithm

- 1: Choose a random pair: $\{\phi_1, \phi_2\} = \{\phi(t_1, x_1, y_1, z_1), \phi(t_2, x_2, y_2, z_2)\}$
- 2: Generate the pair $\{\phi'_1, \phi'_2\} = \{\phi(t'_1, x'_1, y'_1, z'_1), \phi(t'_2, x'_2, y'_2, z'_2)\} = T_i(\phi_1, \phi_2)$
- 3: Draw a random number $c \in [0, 1]$
- 4: **if** $c > P(\phi'_1, \phi'_2)/P(\phi_1, \phi_2)$ **then**
- 5: do **nothing**
- 6: **else**
- 7: perform the exchange $(\phi_1, \phi_2) \mapsto (\phi'_1, \phi'_2)$
- 8: **end if**

Algorithm 2: Genetic Loop

In the notation used below, P is such probability profile and $P(\phi'_i, \phi'_j)$ means that this profile is calculated using the [GA generated] points ϕ'_i and ϕ'_j . It proved useful to chose the mollified importance sampling function as this profile P . Also, note that ϕ_i and ϕ_j , just like ϕ'_i and ϕ'_j , are just particular values of the field ϕ at the sites i and j . A comparison between these two profiles is then performed.

5.3. Euclidean Lattice Simulations

In order to be able to understand better what will happen in the Lorentzian/Minkowski case, a quick and dirty reminder of the Euclidean one is presented below.

In what follows, Φ is the set of all possible field configurations and $||\Phi||$ is its *cardinality* (i.e., the number of elements in the set). For each element of Φ (i.e., for each field configuration) one has to average the observable over all the lattice points (which are computed via Metropolis). Each element in the Markov Chain (generated by the Metropolis MC algorithm) is denoted by $\phi^{[i]}$. To obtain the final result, an average over all field configurations is made. (To make clear that the final goal is the actual lattice computation, the measure is denoted as $[d\phi]$.)

$$\begin{aligned}
\langle O \rangle &= \frac{\int O[\phi] \exp\{-S[\phi]\} [d\phi]}{\int \exp\{-S[\phi]\} [d\phi]}, \\
&= \frac{1}{||\Phi||} \sum_{\phi \in \Phi} \left\{ \frac{\left[\frac{\text{Volume}}{N^d} \right] \cdot \sum_{i=1}^{N^d} O[\phi^{[i]}]}{\left[\frac{\text{Volume}}{N^d} \right] \cdot N^d} \right\}, \\
\therefore \langle O \rangle &= \frac{1}{||\Phi||} \sum_{\phi \in \Phi} \left\{ \frac{1}{N^d} \sum_{i=1}^{N^d} O[\phi^{[i]}] \right\}.
\end{aligned}$$

5.4. Mollified Minkowski Lattice Simulations

In the Minkowski version of the above, the Wick rotation is not performed, therefore we are left with the original form of the exponent,

$$\langle O \rangle = \frac{\int O[\phi] \exp\{i S[\phi]\} [d\phi]}{\int \exp\{i S[\phi]\} [d\phi]}.$$

The above functional will be mollified in order to yield more tractable expressions. The fact that there will be two functional integrations (rather than just one, like above) should not bring many problems. The real question here stems from the fact that the mollifier, η , *mixes* the two of them.

$$\begin{aligned}
\langle O \rangle &= \frac{\int \left\{ \int \eta_\epsilon[\phi - \varphi] O[\varphi] \exp\{i S[\varphi]\} [d\varphi] \right\} [d\phi]}{\int \left\{ \int \eta_\epsilon[\phi - \varphi] \exp\{i S[\varphi]\} [d\varphi] \right\} [d\phi]}; \\
\therefore \langle O \rangle &= \frac{1}{||\Phi||} \frac{1}{||F||} \sum_{\substack{\phi \in \Phi \\ \varphi \in F}} \left\{ \frac{\sum_{i,j=1}^{N^d} \eta_\epsilon[\phi^{[j]} - \varphi^{[i]}] O[\varphi^{[i]}] \exp\{i S[\varphi^{[i]}]\} / W_\epsilon^\Phi[\varphi^{[i]}]}{\sum_{i,j=1}^{N^d} \eta_\epsilon[\phi^{[j]} - \varphi^{[i]}] \exp\{i S[\varphi^{[i]}]\} / W_\epsilon^\Phi[\varphi^{[i]}]} \right\};
\end{aligned} \tag{5.1}$$

where it is understood that the configurations in Φ are chosen with respect to the importance sampling function, $W_\epsilon^\Phi[\phi]$, and the configurations in F are chosen with respect to a *uniform* distribution. This happens because the mollification process (namely the *convolution*) mixes the variables from Φ and F together. Therefore, the only way to implement this “mixing” is by having a uniform distribution for F and implementing the “interaction” with the variables in Φ — via the mollification — explicitly, using η . Note that $||\Phi|| = ||F||$.

Spontaneous Symmetry Breaking

We have been discussing lattice QFT in Lorentzian/Minkowski spacetime because one of the main objectives of this paper is to examine a numerical approach to directly calculate Green's functions in different phases in QFT. This necessitates being able to evaluate path integrals of complex exponentials. The arguments for this have been given elsewhere [9] and will be illustrated in a particular example in the following section, but it is fairly easy to understand why this is the case. The traditional [Feynman] Path Integral formulation of a QFT involves integration of the exponential of the action over every field variable at every space time point. Traditionally, these integrations (assume the action is written in terms of self adjoint fields) range from negative to positive infinity along the real axis. In the limit of small couplings, this form of the [Feynman] Path Integral generates perturbation theory and the results appear (for finite number of expansion terms) to be regular at vanishing coupling. Thus, all expansions that are not regular at vanishing coupling, such as the traditional symmetry breaking expansion of quartic scalar field couplings, are excluded. In order to avoid this restriction and produce all possible solutions of the QFT it is necessary to extend the [Feynman] Path Integral integrations to complex values of the fields in a way consistent with the field equations and reality properties of the theory.

Indeed, as shown in [9] and section 7, the different phases of the theory emerge through the varied boundary conditions consistent with the equations of motion. Equivalently, rather than varying the boundary conditions of equations of motion, the measure of the Path Integral can be changed. In general the number of choice of paths of integration for the path integral correspond to the number of independent solutions of the Schwinger-Dyson differential equations.

If we define a QFT via its [Feynman] Path Integral, all we need to know is the action, $S[\phi, J]$ (for a field ϕ whose source is J), of a given model, for then we can write the generating functional as:

$$\mathcal{Z}[J] = \mathcal{N} \int \exp\{i S[\phi; J]\} \mathcal{D}\phi ,$$

where $\mathcal{N}$ is a normalization constant such that $\mathcal{Z}[J = 0] = 1$. The crucial question that remains unanswered in this approach is: *“How does one properly define the measure*

$\mathcal{D}\phi$?”

The best answer so far (for 4-dimensional systems) says that this can only be done for *free* QFTs, via the use of cylindrical functions [17, 18]. (Note that this only happens in the continuum. In its Lattice formulation, QFT is free from such peculiarities because of the lattice regularization.)

Analogously, defining a QFT via its Schwinger-Dyson equation,

$$\frac{\delta S[-i \frac{\delta}{\delta J}]}{\delta \phi} \mathcal{Z}[J] - J(x) \mathcal{Z}[J] = 0 ;$$

is equivalent to substituting $\phi \mapsto -i \frac{\delta}{\delta J}$ in the action, requiring the equations of motion to be the solutions that extremize it. Note that the Schwinger-Dyson equations are a system of infinitely many [partial] differential equations, one per each point of spacetime.

Therefore, when thinking in terms of differential equations, the boundary conditions are responsible for the phase structure of the theory. Intuitively, the picture that comes to mind is that of a portion of space divided into as many subsets as there are solutions to our equations of motion, such that in each of those regions, the equations of motion satisfy appropriate boundary conditions.

On the other hand, when thinking about the [Feynman] Path Integral, usually it does not seem bothersome that the measure is not properly well-defined. In fact, determining the boundary conditions for the Schwinger-Dyson equations is analogous to determining the measure for the Path Integral. Thus, just like the boundary conditions, the measure is responsible for the phase structure of the theory (in the integral representation of the problem).

For completeness sake, this is how a QFT in different spacetime dimensions, d , behaves:

$d = 0$ For 0-dimensional QFTs — i.e., QFT on a point (the [Feynman] Path Integral degenerates into a simple integral) — there is no such thing as phase transition since there is no such thing as dynamics. However, the phase structure of the theory survives, given by the different boundary conditions (resp. measure) needed in order to determine all the solutions to the equations of motion.

$d = 1$ For 1-dimensional QFTs — i.e., Quantum Mechanics —, again, there is no such thing as phase transitions, for if the theory has 2 different vacua we could take a linear combination of them to be the “real” vacuum state, given that they would be related by tunneling. (For the analogous case in Condensed Matter physics, please refer to [19].)

$d \geq 2$ In this case, phase structure and transition exist; this is the complete scenario. (Note that there is no operator that relates 2 inequivalent θ -vacua of the theory, therefore they belong to different algebras.)

The real lesson to be learned from all of this is that Spontaneous Symmetry Breaking is a phenomena generated by boundary conditions; whether you use them to define

the measure of the Path Integral or to define the Schwinger-Dyson equation is just a matter of personal preference. It is not said anywhere that the limits of the Path Integral have to be real; what we need to have are real observables. In fact, as shown below on section 7, it turns out that in order to have symmetry breaking we need a measure that is not necessarily real: this will enable the computation of all the solutions of a given QFT [9].

Chapter 7

Results

In what follows, the results obtained thus far are presented. In short, they are in 0, 1 (time) and 4 spacetime dimensions.

7.1. Lower Dimensional

Let us start by addressing the 0- and 1-dimensional results. Plainly and simply put, this means that we are solving a simple integral in 0 spacetime dimensions since the [Feynman] Path Integral degenerates into a standard integral and, in the 1-dimensional (time) case, one will be doing Quantum Mechanics.

For the 1-dimensional results, refer to [8]. There, the Quantum Chemistry of the problem is fully treated and addressed. Note, however, that the importance sampling function chosen in [8] is different than the one used in this work: the quantum chemistry was done using (3.5) while we use (3.6).

Below, the 0-dimensional results are summarized and, in order to illustrate the features of the mollifier technique, different properties of those models are made explicit.

7.1.1. Airy Function

The action for this model is given by: $S(x) = x^3/3 + Jx$. This is an interesting model because we can explicitly calculate the partition function and compare it with the results coming from the Mollified Monte Carlo procedure. Moreover, this model has 2 stationary phase points and the results displayed are from the one in the complex plane which is not accessible with normal Monte Carlo. The quantity of interest is:




Figure 2: Plot of the Airy potential, $V(x) = x^3/3$, highlighting the fact that it is not bounded from below.

$$\mathcal{Z}[J] = \frac{\int_{-\infty}^{\infty} \exp\left\{i \frac{x^3}{3} + i J x\right\} dx}{\int_{-\infty}^{\infty} \exp\left\{i \frac{x^3}{3}\right\} dx} \equiv \frac{\text{Ai}(J)}{\text{Ai}(0)}.$$

The [first two] graphs below show the highly oscillatory behavior of the integrand in the partition function: the left one is its the real part, while the right one is its the imaginary part.




Figure 3: Real and Imaginary parts of $e^{i x^3/3 + i J x}$ showing the highly oscillatory nature of the problem.

Below we see the graphs of a particular 2-dimensional slice of the above pair, where $J = -1$: the real (left) and imaginary (right) part of the mollified Airy-integrand (in red, $\epsilon = 0.1$) is in contrast to the non-mollified integrand (green).




Figure 4: Superimposed plots of mollified (red) and non-mollified (green) real and imaginary parts of the integrand $e^{ix^3/3-ix}$, showing the smoothness achieved on the $J = -1$ two-dimensional slice of the previous graphs.

Below we see the graphs of the real (left) and imaginary (right) part of the mollified Airy-integrand for different values of the field x and of the mollification parameter ϵ : analogous to Figure 3 but for the mollified version of the integrand (which corresponds to the red lined plots of Figure 4).




Figure 5: Plots of the real and imaginary parts of the mollified integrand for various values of x and ϵ .

The 2 stationary phase points are given by $x_0 \in \{\pm\sqrt{-J}\}$, where x_0 is the solution of $S'(x_0) = 0$, and the [properly mollified] importance sampling functions are:

$$W_\epsilon^{J \geq 0}[x] = \exp\left\{-\frac{1}{2} \frac{x^2 - J}{\epsilon^2}\right\} + \exp\left\{-\frac{1}{2} \frac{x^2 + J}{\epsilon^2}\right\}; \quad J \geq 0; \quad (7.1)$$

$$W_\epsilon^{J < 0}[x] = \exp\left\{-\frac{1}{2} \left(\frac{x - \sqrt{-J}}{\epsilon}\right)^2\right\} + \exp\left\{-\frac{1}{2} \left(\frac{x + \sqrt{-J}}{\epsilon}\right)^2\right\}; \quad J < 0; \quad (7.2)$$

It is customary to see the source term above called t and identified with time but this is not the approach taken here.

Below we see the graphs of W_ϵ with respect to the ϵ parameter and x , for fixed values of the source: on the left $J = 4$, and on the right $J = -16$,




Figure 6: Plots of (7.1), for $J = 4$, and (7.2), for $J = -16$.




Figure 7: Plots of 2-dimensional slices of (7.1) and (7.2) for various values of ϵ .

It is not difficult to see that there is an optimal value for the parameter ϵ , as shown below in the graph for the information theoretic entropy (see Appendix B for further details): we have the information theoretic entropy for $J \geq 0$ on the left, and $J < 0$ on the right,




Figure 8: Information theoretic entropy for finding the optimal value of the mollification parameter: on the left we have $J \geq 0$, and on the right we have $J < 0$.

Furthermore, it is easy to perform the analytical calculations outlined above (using the Airy functions on the Generating Functional, rather than its mollification) to get the pure evaluation of this model. The results show the mollified answers (given a proper choice of ϵ) are sensationally accurate and show, already in this simple example, that a region not allowable in ordinary Monte Carlo approaches can be computed. In fact, due to the simplicity of this example, modern computers can calculate it straightforwardly, without the need of special tricks to handle the highly oscillatory integrand. The point of this example is twofold: show the precision and the accuracy that the mollifier method can achieve and also to illustrate how this smoothing procedure works.

7.1.2. 0-dimensional ϕ^4 Theory

Also known as ultra-local ϕ^4 , its action is given by: $S[\phi] = \mu \phi^2/2 + g \phi^4/4$. As before, the saddle-point ϕ_0 is such that $S'[\phi_0] = 0$. The graphs for the Action above (positive mass on the left, negative mass on the right) are given by:





Figure 9: ϕ^4 potential for positive and negative values of μ .

Below, we show the highly oscillatory behavior of the integrand of the Partition Function given by the above action, i.e., $I = e^{i(\mu \phi^2/2 + g \phi^4/4)}$, for positive and negative values of the μ parameter:






Figure 10: Plots of the Real and Imaginary parts of the integrand for various values of J and $\mu = 1, g = 1$ and $\mu = -1, g = 1$.

Below, we see the real (left) and imaginary (right) parts of the mollified ϕ^4 -integrand (red) in comparison with their non-mollified counterparts (green):




Figure 11: Superimposed mollified (red) and non-mollified (green) real and imaginary parts of the integrand for $\mu = 1, g = 1, J = -1, \epsilon = 0.1$.

Below, we see the real (left) and imaginary (right) parts of the mollified ϕ^4 -integrand for different values of the field ϕ and of the mollification parameter ϵ :




Figure 12: Plots of the real and imaginary parts of the mollified integrand for various values of the field and the parameter ϵ .

As explained in [9, 12], the three different solutions/phases of this model can be selected via an appropriate choice of boundary conditions: $\Gamma^0 = \mathbb{R}$, $\Gamma^+ = (-\infty, 0) \cup (0, i\infty)$ and $\Gamma^- = (-\infty, 0) \cup (0, -i\infty)$. The leftmost figure below shows those boundaries in the Argand Plane: $\Gamma^0 = \mathbb{R}$ is the green line, $\Gamma^+ = (-\infty, 0) \cup (0, i\infty)$ is the red line while $\Gamma^- = (-\infty, 0) \cup (0, -i\infty)$ is the blue line. The rightmost figure below shows the regions (shaded) of the complex ϕ -plane, ($\phi = \rho e^{i\theta}$), defined by $\cos(4\theta) \geq 0$. Any contour, starting and ending at infinity, within one of these four domains corresponds to a particular solution of the 0-dimensional ϕ^4 .




Figure 13: LEFT: Three different contours that render the Path Integral finite and depict the three distinct solutions obtained in this model. RIGHT: Any contour that starts and finishes inside the same-color shaded regions (following the general directions shown on the left) corresponds to a particular finite solution.

Using the three different contours above we are able to find the three different solutions to the equation of motion of this 0-dimensional QFT. The Schwinger-Dyson equation is given by,

$$-i \left(\mu \frac{\delta}{\delta J} + g \frac{\delta^3}{\delta J^3} \right) \mathcal{Z}[J] = J \mathcal{Z}[J],$$

where we can easily see (again) that there has to be three solutions. These are called *Parabolic Cylinder Functions* and are denoted $U(\mu/g, J)$, $V(\mu/g, J)$, $W(\mu/g, J)$. Those behave in the following manner (see Appendix C):

Regular at $g \rightarrow 0$ Consistent with perturbation theory,

Singular ($\simeq \sqrt{g}$) at $g \rightarrow 0$ Symmetry Breaking,

Singular ($\simeq \exp\{\mu/4g\}$) at $g \rightarrow 0$ Instanton.

Thus,

$$\langle \mathcal{O} \rangle_\epsilon[J] = \frac{\int_{-\infty}^{\infty} \int_{\Gamma^{0,\pm}} \eta_\epsilon(\phi - \varphi) \mathcal{O}[\varphi] \exp\left\{i \frac{\mu}{2} \varphi^2 + i \frac{g}{4} \varphi^4 - i J \varphi\right\} d\varphi d\phi}{\int_{-\infty}^{\infty} \int_{\Gamma^{0,\pm}} \eta_\epsilon(\phi - \varphi) \exp\left\{i \frac{\mu}{2} \varphi^2 + i \frac{g}{4} \varphi^4\right\} d\varphi d\phi}.$$

It should be noted, though, that depending on the chosen contour ($\Gamma^0, \Gamma^+, \Gamma^-$), the above integral will resemble a Fresnel one and, as such, its computation is quite simplified.

For the graphs below, we used the following values for the saddle-points,

$$\phi_0 \in \left\{0, \pm \sqrt{-\mu/2g}\right\},$$

where $S'[\phi_0] = 0$. Therefore, the different importance sampling functions are:

Symmetric Phase Given by the Γ^0 contour, this is the one which is regular in the limit $g \rightarrow 0$ and accessible via perturbation theory,

$$W_\epsilon^0[\phi] = \exp\left\{-\frac{1}{2} \left(\frac{\phi}{\epsilon}\right)^2\right\}; \quad \phi_0 = 0;$$

Solitonic Phase This phase is given by the linear combination of the contours Γ^+ and Γ^- such that $\mu \geq 0$, i.e., this represents the soliton solution,

$$W_\epsilon^{\mu \geq 0}[\phi] = \exp\left\{-\frac{1}{2} \left(\frac{\phi^2 + (\phi_0^+)^2}{\epsilon^2}\right)\right\} + \exp\left\{-\frac{1}{2} \left(\frac{\phi^2 - (\phi_0^-)^2}{\epsilon^2}\right)\right\}; \quad \phi_0^\pm = \pm i \sqrt{\mu/2g}; \quad (\mu \geq 0);$$

Broken-Symmetric Phase This one is given by the linear combination of the contours Γ^+ and Γ^- such that $\mu < 0$, i.e., this represents the solution usually referred to as broken-symmetric,

$$W_\epsilon^{\mu < 0}[\phi] = \exp\left\{-\frac{1}{2} \left(\frac{\phi + \phi_0^+}{\epsilon}\right)^2\right\} + \exp\left\{-\frac{1}{2} \left(\frac{\phi - \phi_0^-}{\epsilon}\right)^2\right\}; \quad \phi_0^\pm = \pm \sqrt{-\mu/2g}; \quad (\mu < 0).$$

Below, we have the graphs of these importance sampling functions:




Figure 14: Plots of the mollified importance sampling functions for various values of the field and ϵ : symmetric ($\phi_0 = 0$), solitonic ($\mu = 1, g = 1$) and broken-symmetric ($\mu = -1, g = 1$).

The graphs below show the entropy for the above importance sampling functions, as discussed in Appendix B:




Figure 15: Information theoretical entropy for finding the optimal ϵ in the symmetric, solitonic and broken-symmetric cases.

7.2. Higher Dimensional

As for the 4-dimensional results [for a Free Scalar QFT] the news is not encouraging. As mentioned in section 3.2, there are severe efficiency constraints in this implementation of the mollification; in section 3.2 three possible choices of solutions to this technical issue were given.

It turns out that, from those three solutions, one is non-efficient while the other two produce analogous results: only the imaginary parts of physically interesting

quantities (generating functional and Green's functions) can be tamed. The real parts continue to show an oscillatory pattern.

Although this is at least disappointing, the solutions obtained are good enough to control the imaginary part's oscillations.

It is our hope that changes in algorithms will allow some version of mollification to work in real world problems.

Part II.

**Solution Space and Moduli Space
Topology**

Introduction

Classical gauge theories are well described through differential geometry, where a gauge field is represented by a connection on a principal fibre bundle P for which the structure group is the symmetry group of the theory; see, for example, [22, 23, 24, 25, 26, 27, 28, 29, 30] and references therein.

The phenomenon of symmetry breaking also has its own geometric formulation [25, 26, 27, 28, 29, 30, 31]: the reduction of the principal bundle P . To this classical picture, the question that arises is that of how does a quantum field theory select its vacua given by the different possible reductions of the principal bundle P . The canonical answer to this question is that radiative corrections (beyond the tree-level approximation) cause the QFT to “jump” from the symmetric phase to the broken-symmetric one(s) [34].

However, [34] already expresses concerns about some issues (e.g., the last paragraph on page 1894 and its continuation on page 1895) that are more explicitly treated in [35]. For more examples about these issues, see [36] (pages 27 and 28 discuss the asymptotic nature of the series expansion, their maximum accuracy and issues regarding Borel summation) and [42].

Loosely speaking, what [35] does, is to compute the Schwinger-Dyson equations (for a given QFT) and note that these differential equations have as many solutions as its order indicates. Moreover, the boundary conditions that determine each of these solutions yield different possible values for the parameters of the theory (mass, coupling constants, etc). Thus, each one of these solutions has its own series expansion (which may or may not be equivalent to the perturbative series) and particular behavior.

The analysis of the boundary conditions of the Schwinger-Dyson equation being responsible for the different solutions of the theory done in [35] is more in the spirit of the self-adjoint extension of the associated Hamiltonian, as done in [43, 44]. Although it may not seem so at first, [35] is equivalent to a latticized approach to QFT and, as such, requires 2 types of infinite limits in order to give rise to its continuum version: the thermodynamic (number of particles in the box) and the volume (size of the box) limit. These limits do not commute and fiddling with them (as done in [35]) is analogous to resumming the perturbative series (as done in [34]).

The present work has the goal of developing a geometrical method that brings to the

foreground these issues of vacuum structure in QFT, its multiple solutions, its moduli space, clearly showing that these different solutions are topologically inequivalent. The fact that different configuration spaces for distinct solutions of the equation of motion of a given Lagrangian have different topologies shows that expansions must be performed separately for each solution of a theory, i.e., each phase has to be regarded and treated as a separate theory.

In this sense, this work streamlines how the parameters of a theory (mass, coupling constants, etc) determine the topology of the vacuum manifold (moduli space) which, combined with the picture presented in [35], gives a prescription of how to examine the solution space of a field theory: The boundary conditions of the Schwinger-Dyson equations determine the parameters of the theory which, in turn, determine its topology.

Classical Field Theory and the Jacobi Metric

A typical action for a scalar field has the form,

$$S[\phi] = \int K(\pi, \pi) - V_\tau(\phi) d^n x ;$$

$$K(\pi, \pi) = \frac{1}{2} \mathbf{g}(\pi, \pi) = \frac{1}{2} g_{\mu\nu} \pi^\mu \pi^\nu .$$

where K is the kinetic quadratic form (bilinear and, in case $\pi \in \mathbb{C}$, hermitian; defining the inner product $\mathbf{g}$) and V_τ is the potential, where the index τ collectively denotes coupling constants, mass terms, etc.

We want to reparameterize our field using the arc-length parameterization such that $\tilde{\mathbf{g}}(\tilde{\pi}, \tilde{\pi}) = \tilde{g}_{\mu\nu} \tilde{\pi}^\mu \tilde{\pi}^\nu = 1$, where $\tilde{\mathbf{g}}$ is the new metric and $\tilde{\pi}^\mu$ is the momentum field redefined in terms of this new parameterization (assuming $V_\tau(\phi)$ contains no derivative couplings). That is, we want to scale our coordinate system in order to obtain a conformal transformation of the metric that normalizes our momentum field.

In fact, there are three possible normalizations, depending on the nature of $\tilde{\pi}$:

$$\tilde{\mathbf{g}}(\tilde{\pi}, \tilde{\pi}) = \begin{cases} 1, & \text{spacelike;} \\ 0, & \text{lightlike;} \\ -1, & \text{timelike.} \end{cases}$$

However, this will not concern us here, once our main focus will be to find the geodesics of $\tilde{\mathbf{g}}$. Therefore, we have a clear distinction among the three foliations of Lorentzian spaces:

$\tilde{\mathbf{g}}(\tilde{\pi}, \tilde{\pi}) > 0$ represents the spacelike-leaf and contains only non-physical objects — after the arc-length reparameterization we will have $\tilde{\mathbf{g}}(\tilde{\pi}, \tilde{\pi}) = 1$;

$\tilde{\mathbf{g}}(\tilde{\pi}, \tilde{\pi}) = 0$ clearly singular, representing the null-leaf where there is no concept of distance, given that all objects here are massless;

$\tilde{g}(\tilde{\pi}, \tilde{\pi}) < 0$ represents the timelike-leaf and contains the physical objects — after the arc-length reparameterization we will have $\tilde{g}(\tilde{\pi}, \tilde{\pi}) = -1$.

In order to find the arc-length parameterization, we will borrow an idea from classical mechanics called *Jacobi's metric* [46]. In the Newtonian setting, Jacobi's metric gives an intrinsic geometry for the configuration space (resp. phase space), where dynamical orbits become geodesics, i.e., it maps every Hamiltonian flow into a geodesic one; therefore, solving the equations of motion implies finding the geodesics of Jacobi's metric and vice-versa. This can be done for any closed non-dissipative system (with total energy E), regardless of the number of degrees of freedom.

Before we proceed any further, let us take a look at a couple of simple examples in order to motivate our coming definition of Jacobi's metric.

9.1. Harmonic Oscillator

Let us consider a model similar to the harmonic oscillator (in $(1+0)$ -dimensions): without further considerations, we simply allow for the analytic continuation of the frequency: $\mu = +\omega^2$ or $\mu = -\omega^2$. Its Lagrangian is given by $L = \frac{1}{2} \dot{q}^2 \mp \frac{\mu}{2} q^2$, where $\mu > 0$, from which we conclude that for a fixed E such that $E = \frac{1}{2} (\dot{q}^2 \pm \mu q^2)$, we have $\dot{q} = \frac{dq}{dt} = \sqrt{2(E \mp \frac{\mu}{2} q^2)}$.

So, in the spirit of what was said above, we want to find a reparameterization of the time coordinate in order to have the normalization $\dot{q} = 1$, where the dot represents a derivative with respect to this new time variable.

Thus,

$$\begin{aligned} \frac{dq}{dt} &= \frac{dq}{ds} \frac{ds}{dt} = \sqrt{2 \left(E \mp \frac{\mu}{2} q^2 \right)} ; \\ \text{if } \frac{ds}{dt} &= \sqrt{2 \left(E \mp \frac{\mu}{2} q^2 \right)} \Rightarrow \frac{dq}{ds} \equiv 1 ; \\ \therefore ds &= \sqrt{2 \left(E \mp \frac{\mu}{2} q^2 \right)} dt ; \\ \Rightarrow \tilde{g}_E &= 2 \left(E \mp \frac{\mu}{2} q^2 \right) g . \end{aligned}$$

As expected, the new metric, $\tilde{g}_E$, is a conformal transformation of the original one, g ; and under this new metric, our original Lagrangian is simply written as,

$$\begin{aligned}
L(q, \dot{q}) &= \frac{1}{2} \dot{q}^2 \mp \frac{\mu}{2} q^2 ; \\
&= \frac{1}{2} g_{ij} \dot{q}^i \dot{q}^j \mp \frac{\mu}{2} q^2 ; \\
&= \frac{1}{2} \mathbf{g}(\dot{q}, \dot{q}) \mp \frac{\mu}{2} q^2 ; \\
\therefore L(q, \dot{q}) &= \tilde{\mathbf{g}}_E(\dot{q}, \dot{q}) .
\end{aligned}$$

Now, as we said above [46], the Euler-Lagrange equations for this conformally transformed Lagrangian are simply the geodesics of the metric $\tilde{\mathbf{g}}_E$. The task of finding the geodesics $\gamma(s)$ of the $\tilde{\mathbf{g}}_E$ metric can be more easily accomplished with the help of the normalization condition, $\tilde{\mathbf{g}}_E\left(\frac{d\gamma}{ds}, \frac{d\gamma}{ds}\right) = 1$, and the initial condition $\gamma(s=0) = 0$, given that, in this fashion, we only need to solve a first order differential equation:

$$\begin{aligned}
&\tilde{\mathbf{g}}_E\left(\frac{d\gamma}{ds}, \frac{d\gamma}{ds}\right) = 1 ; \\
\Rightarrow 2 \left(E \mp \frac{\mu}{2} \gamma^2\right) \mathbf{g}\left(\frac{d\gamma}{ds}, \frac{d\gamma}{ds}\right) &= 1 ; \\
\therefore \gamma'(s) = \frac{d\gamma}{ds} &= \sqrt{\frac{1}{2(E \mp \frac{\mu}{2} \gamma^2)}} ;
\end{aligned}$$

with the initial condition: $\gamma(0) = 0$.

Here we should note that solving for $\tilde{\mathbf{g}}_E(\gamma', \gamma') = 1$ (spacelike-leaf) is analogous to solving $\tilde{\mathbf{g}}_E(\gamma', \gamma') = -1$ (timelike-leaf), once the two cases are symmetrical about the origin.

Analytically solving the equation above yields 2 possible answers, depending on the particular form of the potential ($\mu > 0$ in both cases):

1. $V_+ = +\mu \gamma^2/2$: For the case of a positive pre-factor, we get that $\gamma \sqrt{\mu(2E - \mu \gamma^2)} + 2E \arcsin(\gamma \sqrt{\mu/2E}) - 2s \sqrt{\mu} = 0$; and
2. $V_- = -\mu \gamma^2/2$: For the case of a negative pre-factor, we find that $\gamma \sqrt{\mu(2E + \mu \gamma^2)} + 2E \operatorname{arcsinh}(\gamma \sqrt{\mu/2E}) - 2s \sqrt{\mu} = 0$.

It is clear from the expression for V_+ that $\gamma^2 \leq 2E/\mu$, i.e., the length of the [classical] geodesic is bounded; this does not happen with V_- .

These geodesics, $\gamma_{\pm}$, clearly depend on the parameters E and μ ; therefore, in order to plot $\gamma(s)$, we have to make 2 distinct choices: $E/\mu = 1$ (left plot) and $E/\mu = -1$ (right plot). The last plot comparatively depicts both geodesics.




Figure 1: The top-left-corner plot shows the geodesic for $E/\mu = 1$, while the top-right-corner one depicts it for $E/\mu = -1$. The bottom-center plot superimposes both of them.

This “harmonic oscillator” model already portrays the features that we want to identify in forthcoming applications of this technique: changing the values of the parameters of the potential we can identify 2 distinct types of geodesics which will be related to different solutions on the coming examples.

Note, however, that in general the parameters will not be able to vary freely: they will belong to some specified set; and they will be related to each other, i.e., we will be able to find a function of the parameters that constrains their behavior — this will establish “dualities” among the parameters of a given theory, as will be shown below.

9.2. Cubic Potential

For completeness sake, let us try another example, that of a cubic potential given by: $L = \frac{1}{2} \mathbf{g}(\dot{q}, \dot{q}) + q^3/3$. It’s Jacobi Metric is given by $\tilde{\mathbf{g}}_E = 2(E + q^3/3) \mathbf{g}$ and the equation we need to solve in order to find the geodesics, $\gamma(s)$, of this metric is,

$$\begin{aligned} \tilde{\mathbf{g}}_E(\gamma', \gamma') &= 2(E + q^3/3) \mathbf{g}(\gamma', \gamma') = 1 ; \\ 2(E + q^3/3) (\gamma'(s))^2 &= 1 ; \\ \text{with the initial condition: } \gamma(0) &= 0 . \end{aligned}$$

Once again, we should note that solving for $\tilde{\mathbf{g}}_E(\gamma', \gamma') = -1$ (timelike-leaf) is analogous to what is being done above: the graphs are reflected with respect to each other.

This example is slightly different from the previous one for the following reason: before, it was the relative values of E and μ that determined our two solutions, i.e., one for $E/\mu \geq 0$ and one for $E/\mu < 0$. Now, this cubic theory has *no* free parameters in its potential, therefore the geodesics are labeled by the arbitrary parameter E .

Just as before, the metric will be [artificially] degenerate when $E = V(\gamma)$, which tells us that $\gamma \geq (-3E)^{1/3}$: when $E \geq 0$ the geodesic is allowed to have any length, but when $E < 0$ the geodesic vanishes for some time, after which it starts to grow.

This is very interesting because it essentially says that there is a “lag time” before this solution comes alive: this leaf is non-existent for some “proper time” and then it springs into being quite abruptly.




Figure 2: The top-left-corner plot shows the geodesic for $E = 1$, while the top-right-corner one depicts it for $E = -1$. The bottom-center plot superimposes both of them.

Now that we are done with our examples, we are ready to find the generalization of Jacobi’s metric to the field theoretical setting can be accomplished via the definition of the following conformally transformed metric:

$$\begin{aligned} L &= \frac{1}{2} \mathbf{g}(\pi, \pi) - V_\tau(\phi) ; \\ &\equiv \tilde{\mathbf{g}}_E^\tau(\tilde{\pi}, \tilde{\pi}) ; \end{aligned}$$

where,

$$\tilde{\mathbf{g}}_E^\tau = 2 (E - V_\tau(\phi)) \mathbf{g} ; \quad (9.1)$$

and E (an arbitrary parameter) is the total energy of the system.

Chapter 10

Applications in Quantum Field Theory

We will use the approach in terms of Feynman Path Integrals in order to make things more straightforward. However, we could as well talk in terms of the phase space of a given QFT and its vacuum manifold, i.e., its moduli space.

Starting from the partition function, we have the following ($\hbar = 1$):

$$\begin{aligned}\mathcal{Z}[J] &= \mathcal{N} \int e^{iS[\phi] + i \int J(x) \phi(x) d^d x} \mathcal{D}\phi ; \\ &= \mathcal{N} \int e^{i \int \tilde{\mathbf{g}}_E^T(\tilde{\pi}, \tilde{\pi}) + J(x) \phi(x) d^d x} \mathcal{D}\phi ;\end{aligned}$$

where $\mathcal{N}$ is a normalization constant such that $\mathcal{Z}[J = 0] = 1$, and we have already written the Action in terms of Jacobi's metric.

Now, let us expand the partition function above in terms of its classical part and its quantum fluctuations, i.e., $\phi = \phi_{\text{cl}} + \delta\phi$, and the classical Action is given by $S_{\text{cl}} = S[\phi_{\text{cl}}] = \int_{\mathcal{M}} \tilde{\mathbf{g}}_E^T(\tilde{\pi}_{\text{cl}}, \tilde{\pi}_{\text{cl}})$, where $\mathcal{M}$ is the particular region of spacetime where the integration is performed and $\pi = d\phi$ — for this calculation we are assuming the fluctuations vanish at the boundary, $\delta\phi|_{\partial\mathcal{M}} = 0$. Thus, we have that,

$$\begin{aligned}\mathcal{Z} &= \int \exp\{i S[\phi_{\text{cl}} + \delta\phi]\} \mathcal{D}\phi ; \\ &= \int \exp\left\{i \int_{\mathcal{M}} \tilde{\mathbf{g}}_E^T(\tilde{\pi}_{\text{cl}}, \tilde{\pi}_{\text{cl}}) + \tilde{\mathbf{g}}_E^T(\delta\tilde{\pi}, \delta\tilde{\pi})\right\} \mathcal{D}\phi ; \quad (10.1)\end{aligned}$$

$$= e^{i S_{\text{cl}}} F_{\tau}[\partial\mathcal{M}] ; \quad (10.2)$$

where,

$$F_{\tau}[\partial\mathcal{M}] = \int e^{i S[\delta\phi]} \mathcal{D}(\delta\phi) = \int e^{i \int_{\mathcal{M}} \tilde{\mathbf{g}}_E^T(\delta\tilde{\pi}, \delta\tilde{\pi})} \mathcal{D}(\delta\phi) \approx \frac{1}{\sqrt{\det(-D_{\tau}^2)}} ; \quad (10.3)$$

where $F_{\tau}[\partial\mathcal{M}]$ is the [quantum] fluctuation part of the partition function, the index τ denoting collectively the parameters of the potential (mass, coupling constants,

etc), and $D_\tau^2 = \nabla^\mu \nabla_\mu$ is the Laplace-Beltrami operator constructed from the covariant derivative (∇_μ) associated to the Levi-Civita connection of $\tilde{\mathbf{g}}_E^\tau$. There are two important things to note from (10.3): the fluctuation term can only depend on the parameters of the potential and on the boundary $\partial\mathcal{M}$, i.e., surface terms (that we assumed vanishing); and its structure is analogous to that of the classical part, in that $\tilde{\mathbf{g}}_E^\tau$ is the same on both.

However, the above presumes that there is only one solution to the theory in question. But, the method we are developing is exactly to use Jacobi's metric in order to find and classify *all* of the solutions of the theory given, i.e., we want to be able to use this tool to study the moduli space of the problem at hand. Therefore, we need to generalize the situation above for the case of *many* solutions, which is not a difficult task:

$$\mathcal{Z} = \prod_{\nu=1}^N e^{i S_{\text{cl}}^\nu} F_\tau^\nu[\partial\mathcal{M}] ; \quad (10.4)$$

$$\approx \prod_{\nu=1}^N \frac{e^{i S_{\text{cl}}^\nu}}{\sqrt{\det(-D_\tau^2)}} ; \quad (10.5)$$

where ν counts the number of different solutions (i.e., the number of critical points of $S[\phi]$, including its multiplicity) denoted by ϕ_{cl}^ν , and $S_{\text{cl}}^\nu = S[\phi_{\text{cl}}^\nu]$. Note that, in order to find all of the possible critical points of $S[\phi]$, we need to take into account its τ -dependence, and in doing so we are implicitly assuming that $\tau \in \mathbb{C}$, ensuring we are able to find them all.

There are two major observations to be done at this point:

1. There are two kinds of discontinuities present in the above construction, [37]: first, critical *points* ϕ_{cl}^ν can coalesce, which happens in the complexified catastrophe set in parameter space; second, critical *values* $\Im(i S[\phi_{\text{cl}}^\nu]) = S_{\text{cl}}^\nu$ can coalesce, which happens on the Stokes set in parameter space and corresponds to the appearance or disappearance of a subdominant exponential in a “non-local bifurcation”. These correspond, respectively, to realizing that the partition function is to be taken over *complex* fields (rather than real ones, as is customary), where the contour of integration (rendering the partition function finite, [35]) will ultimately determine the parameter space (these are the Lee-Yang zeros of our theory); and the coalescing of S_{cl}^ν represents the Stokes phenomena of our theory. Both of these will determine the phase structure of our problem (see [35]). Note that it is the contour of integration that connects both of these, once the appropriately chosen range of ϕ will render $e^{i S[\phi]}$ convergent.
2. The asymptotic behavior of the partition function depends *only* on the critical points of S , i.e., on ϕ_{cl}^ν . Thus, when $S[\phi]$ (resp. $V[\phi]$) is a Morse function, i.e, smooth with no degenerate critical points, we can use Morse's lemma to show that the critical points are isolated, keeping in mind that the number of isolated

points is a topological invariant. (It is worth noting this can be generalized in the thermodynamical limit via the Morse-Palais lemma, just as we can relax the condition of non-degeneracy of the critical points via Morse-Bott theory.) Using this, it can be shown that ϕ -space (resp. Phase Space) is a CW-complex with a ν -cell for each critical point of index ν : the fluctuation term contains the Maslov-Morse index (in $\tilde{\mathbf{g}}_E^\tau$ by means of its dependence on $V_\tau[\phi]$) that accounts for the discretization of Path Space and corrects for the thermodynamic limit of the particular solution in question, i.e., every time the denominator in (10.5) vanishes, $F_\tau^\nu[\partial\mathcal{M}]$ passes a singularity in such a way as to ensure the proper phase; the phase factor that arises in this way is nothing but $e^{i\alpha\nu}$, where α is some angle and ν counts the number of zeros (with multiplicity) encountered along the particular path in question — it is called the Maslov-Morse index.

Now, we are ready to employ this machinery in order to draw several important conclusions. Here are them:

1. The partition function, seen as a function of the parameters of the potential, $\mathcal{Z} = \mathcal{Z}[\tau] = \mathcal{Z}[\text{mass, coupling constants}]$, is a *meromorphic* function: it is holomorphic on a subset of $\mathbb{C}$ except for the set of isolated points given by the values of the parameters along Stokes' lines (resp. critical lines of phase transition). We have to keep in mind that the parameter space had to be complexified in order to yield all possible solutions to the theory given; another way to think about this is in terms of the contours that render the partition function finite: changing these contours (looking for all possible ones that make the partition function converge) will affect the allowed values for the parameters, i.e., these contours ultimately determine the parameter space, as done in [35] (see also [40]).
2. Under the [full] elliptic modular group, $\Gamma = \text{SL}(2, \mathbb{Z}) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mid a = 1, b = 0, c = 0, d = 1 \right\}$ (see [38]), the partition function is a *modular function*, i.e., for any $\mathbb{M} \in \Gamma$ we have that $\mathcal{Z}[\mathbb{M}\tau] = \mathcal{Z}[\tau]$, where $\mathbb{M}\tau = \frac{a\tau+b}{c\tau+d}$. (For a fuller appreciation of the importance of this fact, see, e.g., [39].)
3. The Action, written in terms of Jacobi's metric, $S[\phi] = \int_{\mathcal{M}} \tilde{\mathbf{g}}_E(\tilde{\pi}, \tilde{\pi})$, can be thought of as the Morse-theoretic Energy functional, $E[\gamma] = \int \mathbf{g}(\gamma', \gamma')$, for the path γ , where $\gamma' = d\gamma/ds$. Therefore, we can readily say that all of the critical points of $S[\phi]$ are given by minimal [Lagrangian] manifolds, and, just like E'' , $S'' = \delta^2 S / \delta\phi(x) \delta\phi(y)$ is a well defined symmetric bilinear functional. This implies that $S'' = 0$ if, and only if, $\tilde{\pi}$ is a Jacobi field, which, in turn, implies that $\tilde{\pi}|_{\partial\mathcal{M}} = 0$. Therefore, using the split of the partition function (10.4) in terms of a classical part and its quantum fluctuations, we see that the vacuum manifold (moduli space) of the quantum theory is given by its classical minimal Lagrangian manifold with the quantum corrections being given by extensions of it via Jacobi fields. That is, the classical minimal manifold is extended via the gluing of the quantum fluctuation manifold obtained as a solution to the equations of motion coming from $S_{\text{fl}} = \int_{\mathcal{M}} \tilde{\mathbf{g}}_E^\tau(\delta\tilde{\pi}, \delta\tilde{\pi})$. Thus, the quantum corrections are handles attached to the classical solution, without changing the classical topology.

4. Continuing the reasoning above, it is not difficult to see that a phase transition will happen when the quantum corrections (in terms of handle attachments) actually do change the topology of the particular solution in question. In fact, if $V_\tau[\phi]$ crosses a critical point of index ν , the handle to be attached is a ν -cell (i.e., a ν -simplex) — the connection with what has been said before should be straightforward. Analogously, we can say the following: let $\Sigma_\tau^E = V_\tau^{-1}[E] = \{\phi \mid V_\tau[\phi] = E\}$, i.e., Σ_τ^E is the set of all field configurations which have the same potential energy — this manifold is the configuration space (resp. moduli space), an equipotential surface. Thus, the family of equipotentials $\{\Sigma_\tau^E\}_{E \in \mathbb{R}}$ foliates the configuration space (moduli space) in such a way that if Σ_τ^E is $\mathcal{C}^\infty$ -diffeomorphic to $\Sigma_{\bar{\tau}}^{\bar{E}}$ (for two different values of the energy, E and $\bar{E}$) then there is no phase transition. Conversely, a phase transition will be characterized by the existence of a certain critical value E_c such that $\{\Sigma_\tau^E\}_{E < E_c}$ is *not* $\mathcal{C}^\infty$ -diffeomorphic to $\{\Sigma_{\bar{\tau}}^{\bar{E}}\}_{\bar{E} > E_c}$ (note that, upon a phase transition, the parameters of the potential change from τ to $\bar{\tau}$). Further, the topological difference between these two leaves is a ν -cell, where ν is the index of the critical point E_c . Loosely speaking, it can be said that the origin of phase transitions is this change in topology. Therefore, different phases of the theory are topologically inequivalent.
5. Finally, let us note that all of these conclusions we have drawn so far have one last implication, a topological constraint: the Euler Characteristic, χ , computed from $\tilde{\mathbf{g}}_E$ gives the quantization rules [for the energy]. Therefore, our theory can be topologically quantized; in fact, each topologically inequivalent leaf $\{\Sigma_\tau^E\}$ has its own χ_τ^E and, thus, its own quantization rules.

With all of these facts in hand, let us see what is the framework they imply: given a theory, $L = \mathbf{g}(\pi, \pi) + V_\tau[\phi]$, we can readily construct its Jacobi metric, $\tilde{\mathbf{g}}_E^\tau = 2(E - V_\tau) \mathbf{g}$: from this, we can do two things, either compute the geodesics of $\tilde{\mathbf{g}}_E^\tau$ and classify their Jacobi fields in terms of τ , or calculate its Euler Characteristics, χ_τ^E , and find the quantization rules (bearing in mind that they will vary with τ). The different geodesics (corresponding to different Jacobi fields) will label diffeomorphically equivalent foliations $\{\Sigma_\tau^E\}$, while the order of the zeros of $F_\tau^\nu[\partial\mathcal{M}]$ will correspond to the ν -cells associated with a particular phase transition — furthermore, they will be responsible for the Lee-Yang zeros and the Stokes phenomena of the theory. In turn, this gives the partition function a meromorphic and a modular character when seen with respect to τ .

All of this was possible because we extended the parameter space, allowing τ to be complex. This is completely analogous to complexifying the solution space (moduli space) of the theory given, a fact which is made clear in [35] (and employed in lattice calculations in [40]).

Lastly, using τ , it will be possible to construct “dualities” between different phases. However, in general, these dualities will be non-trivial combinations of the parameters of the theory (as opposed to what we found in the simple examples above, where $E/\mu \mapsto -E/\mu$ or $E \mapsto -E$ did the job).

10.1. The $\lambda\phi^4$ Potential

This theory is defined for scalar-valued fields, ϕ , by the Lagrangian $L = \frac{1}{2}(\mathbf{g}(\pi, \pi) - \mu\phi^2 - \frac{\lambda}{2}\phi^4)$, i.e., the potential is given by $V(\phi) = \frac{1}{2}(\mu\phi^2 + \frac{\lambda}{2}\phi^4)$; which is invariant by $\mathbb{Z}_2$ -reflection: $\phi \mapsto -\phi$.

The Jacobi metric for this theory is given by $\tilde{\mathbf{g}}_E^\tau = 2(E - \frac{\mu}{2}\phi^2 - \frac{\lambda}{4}\phi^4)\mathbf{g}$. And solving the normalization condition $\tilde{\mathbf{g}}_E^\tau(\gamma', \gamma') = 1$ — where $\gamma(s)$ is the geodesic we want to compute and $\gamma' = \frac{d\gamma}{ds}$, where s is the arc-length parameter — should give us the geodesic structure of the vacuum manifold (resp. moduli space):

$$\begin{aligned}\tilde{\mathbf{g}}_E^\tau(\gamma', \gamma') &= 1 ; \\ 2(E - \frac{\mu}{2}\gamma^2 - \frac{\lambda}{4}\gamma^4)(\gamma')^2 &= 1 ; \\ \sqrt{E - \frac{\mu}{2}\gamma^2 - \frac{\lambda}{4}\gamma^4} d\gamma &= \frac{1}{2} ds .\end{aligned}\tag{10.6}$$

The full picture presents itself upon a more detailed analysis of $\int (E - \mu\gamma^2/2 - \lambda\gamma^4/4)^{1/2} d\gamma$, which is the [elliptic] integral that needs to be solved in order to find $\gamma(s)$. Therefore, it is useful to consider the polynomial $P(\gamma) = (\gamma^2 - r_1)(\gamma^2 - r_2)$, where $(-\lambda/4)P(\gamma) = E - \mu\gamma^2/2 - \lambda\gamma^4/4$ and $r_{1,2} = -(\mu \pm \sqrt{4E\lambda + \mu^2})/\lambda$. It is the *discriminant* of this polynomial $P(\gamma)$ that will, ultimately, determine the different solutions of the theory: $\Delta = (r_1 - r_2)^2 = \lambda E + \mu^2/4 \gtrless 0$.

As mentioned above, the “dualities” between different phases involves non-trivial combinations of the parameters of the theory. In this case, the dualities are given by the possible values of the discriminant above: $\Delta > 0$, $\Delta = 0$ or $\Delta < 0$.

The two inequalities, $\Delta > 0$ and $\Delta < 0$, are related by the reflection of the τ parameter of this theory, where $\tau = \mu^2/\lambda$, i.e., by the analytic continuation of the mass parameter such that $\mu^2 \mapsto -\mu^2$, which implies that $\tau \mapsto -\tau$ — note that this can be obtained by a modular transformation, $\mathbb{M}\tau = \frac{a\tau+b}{c\tau+d} = -\tau$, such that $b = 0 = c$, $d = 1$ and $a = -1$; further, the analytic continuation of μ is such that $\mu \mapsto \pm i\mu$, or more generically as $\mu \mapsto e^{\pm i\pi\nu/2}\mu$, for odd ν (this is related to the Stokes phenomena discussed previously, where ν selects a certain Riemann sheet for this analytic continuation, which depends on the Maslov-Morse index ν) —; meanwhile, the case $\Delta = 0$ has to be computed separately, once it establishes a fixed value $\tau = -4E$ — note that when $\Delta = 0$ the *resultant* of $P(\gamma)$ and its derivative, $P'(\gamma)$, also vanishes, this resultant being given by $\text{Res}(P, P') = -\frac{\lambda}{4}\Delta$, which says that P and P' have a common root (which happens when $\tau = \frac{\mu^2}{\lambda} = -4E$).

This whole framework is potentially a very interesting result, once discriminants are closely related and contain information about *ramifications* (“branching out” or “branches coming together”) in Number Theory. As we just saw, these ramifications are related to the topological structure of each solution, where the Maslov-Morse index, ν , selects the appropriate Riemann sheet (for the analytic continuation) and the topology of the problem (determined by the attachment of a ν -cell). Therefore, this duality relating the two different solutions, for τ and $-\tau$, is given by a [particular]

modular transformation of τ (resp. analytic continuation of μ), which turns out to be a measure of the ramifications of the problem.

The analytical answers for the geodesic $\gamma(s)$ are found to be given by the following implicit equations:

$\Delta = 0$, $\text{Res}(P, P') = 0$:

$$\frac{1}{3} \gamma^3 + \frac{\mu}{\lambda} \gamma + s = 0 ; \quad (10.7)$$

$\Delta > 0$, $\text{Res}(P, P') < 0$:

$$\begin{aligned} 3s + \gamma \sqrt{(\gamma^2 - r_1)(\gamma^2 - r_2)} + 2r_1 \sqrt{r_2} F\left(\frac{\gamma}{\sqrt{r_1}}; \sqrt{\frac{r_1}{r_2}}\right) - \\ - \frac{2}{3} \frac{\mu}{\lambda} \sqrt{r_2} \left[F\left(\frac{\gamma}{\sqrt{r_1}}; \sqrt{\frac{r_1}{r_2}}\right) - E\left(\frac{\gamma}{\sqrt{r_1}}; \sqrt{\frac{r_1}{r_2}}\right) \right] = 0 ; \end{aligned} \quad (10.8)$$

$\Delta < 0$, $\text{Res}(P, P') > 0$:

$$\begin{aligned} 3s + \gamma \sqrt{(\gamma^2 - r_1)(\gamma^2 - r_2)} + 2r_1 \sqrt{r_2} F\left(\frac{\gamma}{\sqrt{r_1}}; \sqrt{\frac{r_1}{r_2}}\right) - \\ - \frac{2}{3} \frac{\mu}{\lambda} \sqrt{r_2} \left[F\left(\frac{\gamma}{\sqrt{r_1}}; \sqrt{\frac{r_1}{r_2}}\right) - E\left(\frac{\gamma}{\sqrt{r_1}}; \sqrt{\frac{r_1}{r_2}}\right) \right] = 0 . \end{aligned} \quad (10.9)$$

Note that, in the equations above, $F(z; k)$ is the incomplete elliptic integral of the first kind, while $E(z; k)$ is the incomplete elliptic integral of the second kind. Moreover, $r_{1,2}$ are defined (as shown above) as the solutions to the polynomial equation $P(\gamma) = 0$: $r_{1,2} = -(\mu \pm \sqrt{\mu^2 + 4\lambda E})/\lambda$. On top of this, although (10.8) and (10.9) have the same form, they will yield distinct solutions, once the relation given by $\Delta = \lambda E + \mu^2/4$ will either be positive or negative, which, in turn, affects the outcome of $r_{1,2}$ — as already discussed above for $\tau \mapsto -\tau$.

The graphical results are shown below, where $\frac{E}{\lambda} = 1$, $\gamma(0) = 0$ and Δ , respectively, assumes positive ($\Delta > 0$), null ($\Delta = 0$) and negative ($\Delta < 0$) values:




Figure 3: The top-left-corner plot shows the geodesic for $E/\lambda = 1$ and $\Delta > 0$, while the top-right-corner one depicts it for $E/\lambda = 1$ and $\Delta = 0$. The bottom-center plot shows $E/\lambda = 1$ and $\Delta < 0$.

We can clearly see that, for positive Δ (leftmost graph), we have a smooth geodesic $\gamma(s)$ representing the symmetric phase. Then, when Δ vanishes (middle graph), there is a clear change which delimits the 2 different phases of the theory. Moreover, when Δ is negative (rightmost graph), we have the third [broken-symmetric] phase of the theory (which has a finite geodesic).

Gauge Theory Examples

Now, we will consider examples of theories with gauge symmetry (rather than the discrete $\mathbb{Z}_2$ -symmetry of above). As seen in (10.6), taking $\gamma \mapsto -\gamma$ did not change the geodesic structure, i.e., the procedure was covariant (resp. equivariant). This should come as no surprise, once the potential, $V_\tau(\phi)$, is an equivariant function of the fields and the spacetime (base manifold) metric does not depend on the gauge symmetry involved either.

As mentioned before in section 10, we have assumed thus far that the quantum fluctuations vanish at the boundary, $\delta\phi|_{\partial\mathcal{M}} = 0$. This, in turn, implies that the Jacobi fields $\tilde{\pi}$ will satisfy $\tilde{\pi}|_{\partial\mathcal{M}} = 0$. While the fields considered have no internal structure (i.e., gauge symmetry), this is a fairly straightforward constraint, in the sense that its solution is trivial (albeit labeled by τ). However, when gauge symmetry is present, there may be *non-trivial* solutions to these constraints. This is the phenomenon of spontaneous symmetry breaking.

Therefore, following the discussion done in section 10, the following may happen: the quantum corrections, which are handles attached to the classical solution, may now be able to change the topology of the classical solution. So, when a symmetry is reduced from $G \rightarrow H$ (where H is a subgroup of G ; see, e.g., [25, 26, 27, 28, 29, 31]), we start with a Jacobi field which satisfies $\tilde{\pi}_G|_{\partial\mathcal{M}} = 0$ and end up with a Jacobi field who only has H as symmetry, which implies that $\tilde{\pi}_H|_{\partial\mathcal{M}} = 0$ — this means that the initial degrees-of-freedom that combined (respecting the G symmetry) to yield the [initial] constraint $\tilde{\pi}_G|_{\partial\mathcal{M}} = 0$, are not all available now, such that only part of the original symmetry is still respected, yielding $\tilde{\pi}_H|_{\partial\mathcal{M}} = 0$ (the remaining degrees-of-freedom having recombined in non-trivial ways); thus, we end up with a source (or sink) of Jacobi fields that only have H as symmetry.

In this sense, the Atiyah-Singer Index Theorem can be used to measure this variation (i.e., to measure the inequivalent representations of the algebra of observables):

1. If all of the quantum corrections preserve the topology of the classical solution, the topological index does not change, which implies that the analytical index of the differential operator in question also does not change, which, in turn, leaves the vacuum state unchanged.

2. On the other hand, if the quantum fluctuations change the topology of the classical solution (as described above), the topological index will change (following the attachment of the appropriate ν -cell), implying that the analytical index changes as well, which means that the vacuum state changes.

From a different viewpoint, the question can be posed in the following way: Given a certain symmetry breaking connection, how can the topology of the moduli space of the associated Higgs Bundle be studied?

There are some studies in this direction, but no general answers: this is because these types of characterizations are highly model-dependent, i.e., they depend on the particular properties of the connection chosen for the Higgs Bundle in question. For instance, following the discussion above, if the boundary $\partial\mathcal{M}$ is non-existent (i.e., $\mathcal{M}$ is compact), then $\tilde{\pi}|_{\partial\mathcal{M}} = 0$ is trivially satisfied; however, if $\mathcal{M}$ has, e.g., punctures (i.e., a finite set of points omitted), then the connection in question will have a certain ramification structure and the constraint $\tilde{\pi}|_{\partial\mathcal{M}} = 0$ will have non-trivial solutions (see, e.g., [52], and references therein); therefore, the boundary conditions (in the sense of [35, 40]) on the Jacobi fields ultimately determine the structure of the connection (i.e., gauge field).

The examples below are twofold: the first one (Landau-Ginzburg Functional, section 11.1) serves the purpose of showing the equivariance of the method discussed in this work, once the connection (which is $\mathfrak{u}(1)$ -valued) does not undergo symmetry breaking; while the second one (Seiberg-Witten Functional, section 11.2) can be understood as defined for pairs (A, ϕ) , where A is a Hermitian connection (compatible with the holomorphic structure on the bundle in question) and ϕ is a section (of the bundle at hand) — in this sense, one can consider the space of solutions to Hitchin's equations (i.e., the moduli space of the associated Higgs Bundle) given by,

$$F_A + [\phi, \phi^*] = 0 ; \quad (11.1)$$

$$d_A''\phi = 0 ; \quad (11.2)$$

where F_A is the curvature of A and $d_A''\phi$ is the anti-holomorphic part of the covariant derivative of ϕ (compare these with equations (11.11) and (11.12)). In this sense, the different solutions found via (11.13) are a direct statement about the topology of the moduli space of the Higgs Bundle under study.

11.1. The Landau-Ginzburg Functional

To start off, let us consider the case where the base manifold is a compact Riemann surface Σ equipped with a conformal metric and the vector bundle is a Hermitian line bundle $\mathcal{L}$ (i.e., with fiber $\mathbb{C}$ and a Hermitian metric $\langle \cdot, \cdot \rangle$ on the fibers).

The Landau-Ginzburg functional is defined for a section φ and a unitary connection $D_A = d + A$ of $\mathcal{L}$ as ($\sigma \in \mathbb{R}$ is a real scalar),

$$L(\varphi, A) = \int_{\Sigma} |F_A|^2 + |D_A \varphi|^2 + \frac{1}{4} (\sigma - |\varphi|^2)^2.$$

Thus, its Euler-Lagrange equations [of motion] are given by:

$$\begin{aligned} D_A^* D_A \varphi &= \frac{1}{2} (\sigma - |\varphi|^2) \varphi ; \\ D_A^* F_A &= -\operatorname{Re} \langle D_A \varphi, \varphi \rangle ; \end{aligned}$$

where, D_A^* is the dual of D_A , i.e., $D_A^* = - * D_A * = - * (d + A) *$. Note that the equation for F_A (the second one above) is linear in A . Since D_A is a unitary connection, A is a $\mathfrak{u}(1)$ -valued 1-form. This Lie algebra (of the group $U(1)$) will sometimes be identified with $i\mathbb{R}$ — in other words, our A corresponds to $-iA$ in the standard physics literature (where A is real-valued).

Before we go any further, some notational remarks are in order. We decompose the space of 1-forms, Ω^1 , on Σ as $\Omega^1 = \Omega^{1,0} \oplus \Omega^{0,1}$, with $\Omega^{1,0}$ spanned by 1-forms of the type dz and $\Omega^{0,1}$ by 1-forms of the type $d\bar{z}$. Here $z = x + iy$ is a local conformal parameter on Σ and $\bar{z} = x - iy$. Therefore, $dz = dx + i dy$, $d\bar{z} = dx - i dy$, $\partial_z = \frac{1}{2} (\partial_x - i \partial_y)$ and $\partial_{\bar{z}} = \frac{1}{2} (\partial_x + i \partial_y)$. Furthermore, if ∂_x and ∂_y are an orthonormal basis of the tangent space of Σ at a given point, we have that $\langle dz, dz \rangle = 2$, $\langle d\bar{z}, d\bar{z} \rangle = 2$ and $\langle dz, d\bar{z} \rangle = 0$. Given that the decomposition $\Omega^1 = \Omega^{1,0} \oplus \Omega^{0,1}$ is orthogonal, we may also decompose D_A accordingly: $D_A = \partial_A + \bar{\partial}_A$, where $\partial_A \varphi \in \Omega^{1,0}(\mathcal{L})$ and $\bar{\partial}_A \varphi \in \Omega^{0,1}(\mathcal{L})$ for all sections φ of $\mathcal{L}$ (holomorphic, $\bar{\partial}_A f(z, \bar{z}) = 0 \Leftrightarrow f(z, \bar{z}) = f(z)$, and anti-holomorphic, $\partial_A f(z, \bar{z}) = 0 \Leftrightarrow f(z, \bar{z}) = f(\bar{z})$, parts; i.e., the space of 1-forms and the space of connections is decomposable into a direct sum of its holomorphic and anti-holomorphic parts: $\partial_A = \partial + A^{1,0}$ and $\bar{\partial}_A = \bar{\partial} + A^{0,1}$; while the exterior derivative is given by $d = \partial + \bar{\partial}$). As expected, we have that $\partial_A \bar{\partial}_A = 0 = \bar{\partial}_A \partial_A$ and $F_A = -(\partial_A \bar{\partial}_A + \bar{\partial}_A \partial_A)$.

It is not difficult to show [50] that:

$$\begin{aligned} L(\varphi, A) &= \int_{\Sigma} |F_A|^2 + |D_A \varphi|^2 + \frac{1}{4} (\sigma - |\varphi|^2)^2 ; \\ &= 2\pi \deg(\mathcal{L}) + \int_{\Sigma} 2 |\bar{\partial}_A \varphi|^2 + \left(*(-i F_A) - \frac{1}{2} (\sigma - |\varphi|^2) \right)^2 ; \end{aligned}$$

$$\text{where } \deg(\mathcal{L}) = c_1(\mathcal{L}) = \frac{i}{2\pi} \operatorname{tr}(F_A) ;$$

i.e., the degree of the line bundle is given by the 1st Chern class.

Therefore, a very useful consequence from the above is that if $\deg(\mathcal{L}) \geq 0$ the lowest possible value for $L(\varphi, A)$ is realized if φ and A satisfy

$$\bar{\partial}_A \varphi = 0 ; \tag{11.3}$$

$$*(i F_A) = \frac{1}{2} (\sigma - |\varphi|^2) . \tag{11.4}$$

These equations are just the expression of the self-duality of the Landau-Ginzburg functional. If $\deg(\mathcal{L}) < 0$, then these equations cannot have any solution, thus one has to consider the self-duality equations arising from the Landau-Ginzburg functional where the term $+2\pi \deg(\mathcal{L})$ is substituted by $-2\pi \deg(\mathcal{L})$. Therefore, without loss of generality, we shall assume $\deg(\mathcal{L}) \geq 0$. A necessary condition for the solvability of (11.4) is that,

$$\begin{aligned} 2\pi \deg(\mathcal{L}) &= \int i F_A = \frac{1}{2} \int_{\Sigma} (\sigma - |\varphi|^2) \leq \frac{\sigma}{2} \text{Area}(\Sigma) ; \\ \therefore \sigma &\geq \frac{4\pi \deg(\mathcal{L})}{\text{Area}(\Sigma)} ; \end{aligned} \quad (11.5)$$

and the equality only occurs if, and only if, $\varphi \equiv 0$.

The following result is useful when studying the solutions of the above functionals [50]: *Let Σ be a compact Riemann surface with a conformal metric and $\mathcal{L}$ as before. For any solution of (11.3), we have that $|\varphi| \leq \sigma$ on Σ .* That is, this maximum principle states that the amplitude of the field cannot exceed the height of the potential (see the first plot below).


Let us now construct the Jacobi metric for this potential and find its possible geodesics:

$$\begin{aligned} \tilde{g}_E &= 2(E - V_{\sigma}(\gamma)) \mathbf{g} ; \\ &= 2 \left(E + \frac{1}{4} (\sigma - |\gamma|^2)^2 \right) \mathbf{g} . \end{aligned}$$

In a complete analogy to what was previously done, we consider the $P(\gamma) = E - V(\gamma)$ polynomial. However, in order to make this analysis clearer, let us, in fact, use a slightly different polynomial given by $P'(\gamma) = 4(P(\gamma) - E)$. It is straightforward to see that $P'(\gamma) = (|\gamma|^2 - \sigma)(|\gamma|^2 - \sigma)$, which means that σ is the only root of $P'(\gamma)$, with double multiplicity (the two roots coalesce into one).

This situation implies that the discriminant of $P'(\gamma)$ vanishes, i.e., $\Delta = 0$, and the different phases of the theory are labelled by $\sigma = 0$ and $\sigma > 0$.

The plots below have, respectively, the following values for σ : 0.0, 3.0, 5.0, 7.0, 11.0 and 31.0. As they show, when $\sigma = 0$ we have $c_1(\mathcal{L}) = \deg(\mathcal{L}) = \frac{i}{2\pi} \text{tr}(F_A) = 0$ and its geodesic has a clear character which is quite different otherwise:






Figure 4: The plots above show the clear distinction between the $\sigma = 0$ solution and the $\sigma > 0$ ones; the first plot, on the upper left corner, has $\sigma = 0$, while the other ones have differing positive values for σ .

As a last remark on this model, note it can be thought as defined by a pair (A, φ) , where A is a unitary connection compatible with the holomorphic structure of the bundle in question, and φ is a [global] section of this bundle. In this sense, Hitchin's self-duality equations become,

$$\bar{\partial}_A \varphi = 0 ; \quad (11.6)$$

$$F_A + [\varphi, \varphi^*] = 0 . \quad (11.7)$$

Compare these with (11.3) and (11.4).

In this sense, the solutions found above are a direct statement about the topology of the Higgs Bundle in question.

11.2. The Seiberg-Witten Functional

In this case, the base manifold, $\mathcal{M}$, is a compact, oriented, 4-dimensional Riemannian manifold endowed with a spin^c structure, i.e., a spin^c manifold. The determinant line of this spin^c structure will be denoted by $\mathcal{L}$ and the Dirac operator determined by a unitary connection A on $\mathcal{L}$ will be denoted by $\mathcal{D}_A$. Recalling the half spin bundle $\mathcal{S}^\pm$ defined by the spin^c structure, we see that $\mathcal{D}_A$ maps sections of $\mathcal{S}^\pm$ into sections of $\mathcal{S}^\mp$ [47, 48, 49, 50].

In this fashion the Seiberg-Witten functional for a unitary connection A on $\mathcal{L}$ and a section φ of $\mathcal{S}^+$ is given by,

$$SW[\varphi, A] = \int_{\mathcal{M}} |\nabla_A \varphi|^2 + |F_A^+|^2 + \frac{R}{4} |\varphi|^2 + \frac{1}{8} |\varphi|^4 ; \quad (11.8)$$

where ∇_A is the spin^c connection induced by A and the Levi-Civita connection of $\mathcal{M}$, F_A^+ is the self-dual part of the curvature of A and R is the scalar curvature of $\mathcal{M}$. Its Euler-Lagrange equations are given by,

$$\nabla_A^* \nabla_A \varphi = - \left(\frac{R}{4} + \frac{1}{4} |\varphi|^2 \right) \varphi ; \quad (11.9)$$

$$d^* F_A^+ = - \text{Re} \langle \nabla_A \varphi, \varphi \rangle . \quad (11.10)$$

Using a spin frame, it is not difficult [50] to show that the Seiberg-Witten functional can be written in the following form:

$$SW[\varphi, A] = \int_{\mathcal{M}} |\mathcal{D}_A \varphi|^2 + \left| F_A^+ - \frac{1}{4} \langle e_j \cdot e_k \cdot \varphi, \varphi \rangle e^j \wedge e^k \right|^2 ;$$

where e^j are 1-forms dual to the tangent vectors e_j , $e^j(e_k) = \delta_k^j$; $j, k = 1, \dots, 4$. As a corollary of the above, the lowest possible value of the Seiberg-Witten functional is achieved if φ and A are solutions of the *Seiberg-Witten equations*:

$$\mathcal{D}_A \varphi = 0 ; \tag{11.11}$$

$$F_A^+ = \frac{1}{4} \langle e_j \cdot e_k \cdot \varphi, \varphi \rangle e^j \wedge e^k . \tag{11.12}$$

Thus, self-duality is at work yet again: the absolute minima of the Seiberg-Witten functional satisfy not only the second order equations (11.9) and (11.10), but also the first order Seiberg-Witten equations (11.11) and (11.12).

Although our discussion of the Seiberg-Witten functional, so far, has mirrored our discussion of the Landau-Ginzburg one, the parameter σ on the latter has had no analogue in the former. This can be accomplished with the introduction of a 2-form μ and the consideration of the perturbed functional,

$$\begin{aligned} SW_\mu[\varphi, A] &= \int_{\mathcal{M}} |\mathcal{D}_A \varphi|^2 + \left| F_A^+ - \frac{1}{4} \langle e_j \cdot e_k \cdot \varphi, \varphi \rangle e^j \wedge e^k + \mu \right|^2 ; \\ &= \int_{\mathcal{M}} |\nabla_A \varphi|^2 + |F_A^+|^2 + \frac{R}{4} |\varphi|^2 + \left| \mu - \frac{1}{4} \langle e_j \cdot e_k \cdot \varphi, \varphi \rangle e^j \wedge e^k \right|^2 + 2 \langle F_A^+, \mu \rangle . \end{aligned}$$

Their corresponding first order equations of motion are,

$$\mathcal{D}_A \varphi = 0$$

$$F_A^+ = \frac{1}{4} \langle e_j \cdot e_k \cdot \varphi, \varphi \rangle e^j \wedge e^k - \mu .$$

If we assume that μ is closed and self-dual, then we see that $\langle F_A, \mu \rangle = \langle F_A^+, \mu \rangle$, once $\langle F_A^-, \mu \rangle = 0$ due to the orthogonality between anti-self-dual and self-dual forms. Thus, given that F_A represents the first Chern class $c_1(\mathcal{L})$ of the line bundle $\mathcal{L}$, and we assumed μ to be closed (hence it represents a cohomology class $[\mu]$), the integral

$$\int_{\mathcal{M}} \langle F_A, \mu \rangle ,$$

does not depend on the connection A , thus representing a topological invariant, denoted by $(c_1(\mathcal{L}) \wedge [\mu])[\mathcal{M}]$.

Just as before, we also have a maximum principle: *For any solution φ of (11.9) —*

in particular, for any solution of (11.11) — on a compact 4-dimensional Riemannian manifold, we have that,

$$\max_{\mathcal{M}} |\varphi|^2 \leq \max_{x \in \mathcal{M}} (-R(x), 0) .$$

As a direct consequence of this, if the compact, oriented, Riemannian Spin^c manifold $\mathcal{M}$ has nonnegative scalar curvature, the only possible solution of the Seiberg-Witten equations is,

$$\varphi \equiv 0 ; \quad F_A^+ \equiv 0 .$$

The Jacobi metric for this Seiberg-Witten model is given by

$$\tilde{g}_E = 2 \left(E + \frac{R}{4} |\gamma|^2 + \frac{1}{8} |\gamma|^4 \right) g ; \quad (11.13)$$

for a geodesic γ ; and, just like before, although $P(\gamma) = E - V(\gamma) = E + R|\gamma|^2/4 + |\gamma|^4/8$, let us consider $P'(\gamma) = 8(P(\gamma) - E) = (|\gamma|^2 - 0)(|\gamma|^2 + 2R)$. It is then clear that when $R = 0$ the discriminant vanishes ($\Delta = 0$) and the 2 roots merge into 1, what constitutes one of the phases of the theory. When $R \neq 0$, the discriminant is either $\Delta > 0$ ($R > 0$) or $\Delta < 0$ ($R < 0$), which accounts for the other 2 phases of the theory.

The plots below were obtained with the choice of $E = 1.0$ and R respectively equal to 0.0, 3.0, 7.0, -3.0, -5.0, -7.0.




Figure 5: The plot in the upper left corner depicts the solution for $R = 0$, while the plots in the upper center and upper right corner show solutions for $R > 0$. The plots in the lower row picture the solutions for $R < 0$.

Chapter 12

Conclusions

Using a conformal transformation that amounts to finding the arc-length reparameterization of the given problem, we were able to construct a new metric, called Jacobi metric, such that its geodesic equation is equivalent to the original equations of motion.

Then, we separated the given problem into its classical part and quantum fluctuations and, by realizing that the action written in terms of the Jacobi metric is the so-called “energy” in Morse Theory (resp. Morse-Bott theory), we see can reinterpret this as a “topological expansion”, in the sense that the quantum fluctuations are handles attached to the classical solution.

Further, when these handles change the original (classical) topology, via the gluing of an appropriate ν -cell, there is a phase transition, in the sense that we move from one solution [of our QFT] to another. This can be measured via the use of an Index Theorem: once it relates the analytical index of the differential operator in question with the topological index of the manifold under study, we clearly see that if the topological index changes (because of a certain handle attachment), the analytical index must change as well; which means that the zero modes, the solutions of the equations of motion, have to change.

Therefore, by studying the geodesics obtained from the Jacobi metric with respect to its parameters, we can compute the topological index (via generalizations of the Gauss-Bonnet theorem, using the Jacobi metric we derived before) and thus compute the zero-mode solutions with respect to the different values of the parameters. In this way, we are able to classify all possible solutions to the QFT in question, keeping in mind that the quantum corrections are given by handle attachments that may or may not (depending on the values of the parameters) change the topology of the classical solutions.

To this picture, we add Lee-Yang zeros and Stokes phenomena, in order to obtain a more robust view of what is at stake: the Lee-Yang zeros accumulate along Stokes lines and pinch the parameter space, creating different regions of “allowed values” for the parameters of the theory (mass, coupling constants, etc). This represents different sectors of theory, different phases of the theory. And we showed that these are topologically inequivalent.

In this fashion, the partition function is meromorphic, once it is singular along the

Stokes' lines. However, it has various degrees of modular symmetry, depending on the particulars of the theory in question. And, in this sense, different sectors of the theory are related to each other [by an appropriate modular transformation].

In turn, this implies that different solutions of the theory are related to each other, a fact that we dubbed “duality”. Ultimately, these dualities are determined by the actual values of the parameters (and how they compose in order to create the modular symmetry in question), which are determined by the boundary conditions of the Schwinger-Dyson equations (as explained earlier).

Future work will focus on D-modules and dimensional construction of 0-dimensional theories (which can be completely solved). We also intend to generalize the cubic potential showed earlier to matrix- and Lie-algebra-valued fields [53], analytically solving the its 0-dimensional counterpart and dimensionally constructing it (via D-modules): it seems plausible that such an extension might be related to current developments in three-dimensional gravity, [39].

Part III.

Three-Dimensional Gravity and Airy Functions

Chapter 13

Introduction

Recently, new developments have shone new light on the problem of three (1 temporal, 2 spatial) dimensional gravity, [39]. Loosely speaking, here is the gist of the matter.

Classically, 3-dimensional gravity (plus a cosmological constant) is given by the action,

$$S = \frac{1}{16\pi G} \int \left(R + \frac{2}{\ell^2} \right) \sqrt{g} d^3x .$$

It is not difficult to see that the solutions to the above are all locally equivalent and there are no gravitational waves. Thus, naïvely, one may think that this problem is tractable quantum mechanically.

However, on second thought, power counting tells us that G has dimensions of length, rendering the theory unrenormalizable. This leads to the conclusion that the quantum theory does not exist. But this is premature as well, once divergences in perturbation theory can be removed by field redefinitions ($g_{\mu\nu} \mapsto g_{\mu\nu} + a R_{\mu\nu} + \dots$) and a renormalization of ℓ^2 .

Therefore, a slightly different venue of attack must be drawn in order to resolve the conflicts above. What is customarily done is to write the above Action as a gauge theory, which will lead us towards a Chern-Simons theory, in the following way: write the metric in terms of its Cartan connection, $g_{\mu\nu} = \eta_{\mu\nu} e_\mu^i e_\nu^j$, known as the “vielbein” (triad). From this, we write the spin connection, which is an $\mathfrak{so}(2, 1)$ -valued 1-form: $\omega_\mu^{ab} = e_\mu^a \partial_\mu e^{\nu b} + e_\nu^a e^{\sigma b} \Gamma_{\sigma\mu}^\nu$.

At this point, we are ready to combine the vielbein and the spin connection into an $\mathfrak{so}(2, 2)$ -valued connection, A :

$$A = \begin{pmatrix} \omega & \frac{e}{\ell} \\ -\frac{e}{\ell} & 0 \end{pmatrix} .$$

The Chern-Simons action, which is the gauge theoretical version of this problem, is thus given by,

$$S = \frac{k}{4\pi} \int \text{tr} \left(A \wedge dA + \frac{2}{3} A \wedge A \wedge A \right) .$$

Now, we can try and tackle this problem perturbatively. What we see is that, once there is no appearance of the metric tensor in the action above, there are no local densities in this problem, so there are no counterterm contributions in this sense.

Further, we realize that the cosmological constant plays the role of the structure constant of the gauge group. Therefore, it is not renormalizable.

This Chern-Simons description is valid only if the vielbein is invertible, which is true for classical solutions; perturbation theory will not change this.

Finally, this theory is finite and renormalizable by power counting, once there are no local counterterms. Therefore, the quantum theory seems to exist.

However, let us make some non-perturbative considerations:

- There is some unclear relationship between non-invertible vielbeins and the above picture for 3-dim gravity: classical solutions analogous to $A = \omega = e = 0$, which are clearly non-geometrical, must be included in a quantized theory of 3-dim gravity.
- The equivalence between diffeomorphisms and gauge transformations is not straightforward, in the sense that gauge transformations are continuously connected to the identity, while more general diffeomorphisms also play a role in 3-dim gravity. There is no “natural” way to address this situation.
- Further, the above picture in terms of a Chern-Simons theory does not require a sum over topologies, a basic ingredient of a theory of quantum gravity. There is no *a priori* reason to add this to the picture described so far.
- Lastly, for negative values of the cosmological constant, there is a whole class of classical solutions known as BTZ black holes (which make their appearance in the AdS/CFT correspondence). This clearly shows that the theory cannot be “trivial”.

Therefore, we can summarize the above results in terms of the cosmological constant, Λ , in the following fashion:

- $\Lambda > 0$: it is not possible to define any set of observables that can be intrinsically measured.
- $\Lambda = 0$: this theory has no gravitons and no black holes, thus it has no S-matrix (no degrees-of-freedom).
- $\Lambda < 0$: this theory contains BTZ black holes as degrees-of-freedom; therefore its dual 2-dim CFT can be computed via the AdS/CFT duality.

So, at this point, the plan shifts to being that of finding the 2-dim CFT dual (via AdS/CFT) to the Chern-Simons theory above, with negative cosmological constant; for we know that there are degrees-of-freedom in this theory, given by the BTZ black holes.

In this sense, the Euclidean partition function can be written as,

$$\mathcal{Z}[\beta, \theta] = \text{tr} \exp\{-\beta H - i \theta J\};$$

where H is the Hamiltonian, J is the angular momentum (rotation of the asymptotic AdS_3), and β is the imaginary time. This integral is taken over 3-geometries conformal at infinity to a 2-torus with modular parameter $\tau = \theta/2\pi + i\beta$.

However, there are two problems with this picture:

1. The sum of known contributions is not physically sensible because it cannot be written as $\text{tr} \exp\{-\beta H - i \theta J\}$ such that $[H, J] = 0$.
2. This partition function is not convergent, once the action is not bounded from below.

To remedy these, we note the following [39]:

- Real saddle-points do not account for the whole theory.
- Complex saddle-points are needed, which lead us towards a holomorphic factorized partition function consistent with an interpretation as $\text{tr} \exp\{-\beta H - i \theta J\}$.
- Classical geometry: in the semiclassical limit, where $G \rightarrow 0$ with fixed AdS radius ℓ , we obtain *complex* β and θ — justifying the complex saddle-points above.
- Finally, the Hawking-Page phase transition with respect to β , between a thermal gas and a black hole, amounts to being a condensation, on the phase boundary, of the Lee-Yang zeros of the partition function.

It is at this point that we make contact with what was said in Part II of this thesis, and also in [12, 13, 40, 41]: the parameter space (i.e., the (β, θ) -space) must be complexified in order to account for all possible solutions of the theory — this implies that *complex geometries* must be considered in order to make sense of this theory.

Furthermore, all ingredients already described in Part II make their appearance here as well: Lee-Yang zeros accumulate along Stokes' lines in order to account for the Hawking-Page phase transition mentioned, and this corresponds to a topological transition between the two phases. Also, in this case, the partition function has modular symmetry, $\mathcal{Z}[\tau] = \mathcal{Z}[\mathbb{M}\tau]$, where $\mathbb{M} \in \text{SL}(2, \mathbb{C})/\mathbb{Z}_2$.

So, let us see what we can say about this in the next chapters.

Chapter 14

Extended Airy Functions

The first thing we notice is that the Airy functions (Ai and Bi) can be extended from scalar-valued to matrix- and lie-algebra-valued functions, as done in [53].

In this sense, we end up with an integral representation of the form,

$$\text{Ai}(\mathbf{J}) = \int_{\mathcal{V}} e^{i \text{tr}(\frac{\Phi^3}{3} - \mathbf{J} \Phi)} d\Phi ; \quad (14.1)$$

where $\mathcal{V}$ can be either the space of $n \times n$ Hermitian matrices (for the case of a matrix-valued Airy function) or, in general, a vector space that contains the representation of the Lie algebra in question (for the case of a Lie-algebra-valued Airy function); and $\mathbf{J} \in \mathcal{V}$.

This integral does not exist in the sense of Lebesgue, but it is well-defined as a distribution. The formal similarity of these extensions of the Airy function to the scalar-valued Airy function accounts for its name.

In this sense, just as remarked before in chapter 9.2 of Part II, an integral that, *a priori*, looks ill-defined (for it is unbounded from below), does indeed have a finite solution and non-trivial content, given by the Airy function (and its respective extensions). However, this extensions work for more general polynomials of Φ , as long as these have the so-called *Airy property*, [53]. In particular, the differential equations associated with this kind of integral representations is treated by Harish-Chandra's study of invariant differential operators and integrals on semisimple Lie algebras, [54]. Unfortunately, the study of the connections between Harish-Chandra theory, [54], and the Schwinger-Dyson equations of these models is beyond the scope of this work.

It is worth remarking that these extensions of Φ and $\mathbf{J}$ are completely analogous to what we did before when we complexified the ϕ -space. In fact, this is a generalization of that idea: you can extend from $\mathbb{R}$ -valued to $\mathbb{C}$ -valued scalars, but also to more complicated fields, as matrix- or Lie-algebra-valued ones.

Once again, as observed before, the allowed values of $\mathbf{J}$ are determined by the particular contours that render the integral representation above finite, and there will be a certain “duality” between the different solutions determined by the distinct set of allowed values (which is, in turn, determined by the particular contour in question).

Therefore, summarizing, we have managed to generalize the original cubic potential

in different ways: from $\mathbb{R}$ -valued to $\mathbb{C}$ -valued scalars, and from scalars to matrix- and Lie-algebra-valued fields, via equation (14.1). Now, two points are important to note:

1. Equation (14.1) is just a particular case of a more general one, given by [53]:

$$\mathcal{Z}_p(\mathbf{J}) = \int_{\mathcal{V}} e^{i(p(\Phi) - \text{tr}(\mathbf{J}\Phi))} d\Phi ; \quad (14.2)$$

where $p(\Phi)$ is a G -invariant polynomial on $\mathcal{V}$, and G is the Lie group in question. Thus, the Airy case is recovered when we specialize to the case of $p(\Phi) = \Phi^3/3$; but other generalizations are possible, e.g., $p(\Phi) = \text{tr} \Phi^4/4$, or $p(\Phi) = \text{tr}(\mathbf{A} \Phi^2/2 + \mathbf{B} \Phi^4/4)$, where $\mathbf{A}, \mathbf{B} \in \mathcal{V}$.

2. The equation above can be understood as defining a matrix model, where the general connected matrix-model partition function is given by,

$$e^{\mathcal{Z}} = \int e^{-\text{tr}(V(\mathbf{M}))} d\mathbf{M} ;$$

where $\mathbf{M}$ is an $n \times n$ -matrix, and $V(\mathbf{M})$ is the Potential for the model; the integral is taken over the space called $\mathcal{V}$ above.

At this point, we note that in spite of all of these generalizations we have done so far, we are still talking about a 0-dimensional problem. This is no surprise, for we started with a 0-dim problem and changed it in many different ways, but we never added any dimension(s) to it. This is the next step, to construct the dimensions.

Chapter 15

Dimensional Construction Sketch

At this stage, the question poses that itself is: “Given the generalization (14.2), how does one go about engineering the dimensions in order to extend this problem from 0-dimensions to d -dimensions?”

This is a very hard problem and, although it is outside of the scope of this work to tackle it, we will give a very rough sketch of what should be done.

The whole problem of using (14.2) in order to engineer the dimensions from first principles is that two limiting processes are involved: the limit where the size of each dimension tends to infinite (dubbed “infinite volume limit”), and the limit where the number of points in each dimension tends to infinite (called “thermodynamical limit”). What we want to do is to “grow” a space whose natural differential operator is given, in some sense, by a combination of a derivative(s) and $p'(\Phi)$.

In our case (Chern-Simons), the Schwinger-Dyson equations of motion are symbolically given by,

$$\begin{aligned} (-i \, d\mathbf{A} - \mathbf{A} \wedge \mathbf{A}) \, \mathbb{Z}[\mathbf{J}] &= \mathbf{J} ; \\ (-i \, d(\delta\mathbf{J}) - \delta\mathbf{J} \wedge \delta\mathbf{J}) \, \mathbb{Z}[\mathbf{J}] &= \mathbf{J} ; \end{aligned}$$

where $\mathbf{A} \mapsto -i \, \delta\mathbf{J}$, and $\mathbf{A}, \mathbf{J} \in \mathfrak{so}(2, 2)$. In this case, the operator given by $(-i \, d - \mathbf{A} \wedge \mathbf{A})$ is the natural differential operator under study, and it is constructed out of the polynomial $p'(\Phi) = \text{tr} \, \mathbf{A}^2 = \mathbf{A} \wedge \mathbf{A}$, with the derivative operator being given by $d\mathbf{A} = \text{tr}(d\mathbf{A})$.

In some sense, the tool that allows us to engineer however many dimensions as we want is called *D-Module*. By definition, a *D-module* is a module¹ over a ring D of differential operators. The methods of *D-module* theory have always been drawn from sheaf theory and other techniques with inspiration from the work of Alexander Grothendieck in algebraic geometry. The approach is global in character, and differs from the functional analysis techniques traditionally used to study differential operators. The strongest results are obtained for over-determined systems (holonomic systems), and on the characteristic variety cut out by the symbols, in the good case for which it

¹A module is a generalization of a vector space over a field $\mathbb{K}$ when $\mathbb{K}$ is replaced by a ring.

is a Lagrangian submanifold of the cotangent bundle of maximal dimension (involutive systems).

For more about the use of D -modules as partition functions, we refer the reader to [60] (and references therein). In particular, for our problem we want to compute the so-called “Airy D -Module” — to learn more about such problems, please refer to [61] (and references therein).

As a final note, we remark on the properties that will be determined by the two contours that define (14.1). As we already know, these contours (resp. the boundary conditions for the Schwinger-Dyson equations) ultimately define the ranges that the parameters (mass, coupling constants, etc) of the theory can attain. More generally, when talking about gauge theories, we should realize that, mathematically, the coupling constants are the parameters describing the scalar product on the particular Lie algebra in question, while, physically, the coupling constants appear in the structure constants [defining the Lie algebra under study] and is interpreted as measuring the “intensity” of the interaction between fields, [62].

This means that different contours will determine equivalence classes of structure constants: coupling constants in the same equivalence class determine the same Lie algebra, while coupling constants in different equivalence classes will determine distinct Lie algebras.

Therefore, the two contours that yield the A_i and B_i functions and its respective extensions (as defined previously) will ultimately determine different Lie algebras, i.e., two distinct theories.

This is not much of a surprise when we think in terms of the results already derived in Part II, where the geodesics of the cubic potential (chapter 9.2 of Part II) were shown to be significantly different (thus, the group manifolds in question have to be just as distinct).

In particular, in this context of 3-dimensional gravity, one of the solutions is particularly interesting, for it vanishes for a certain time, after which it suddenly and abruptly starts to grow. It begs the question: does it really mean that, in this particular solution, (2+1)-spacetime is non-existent for some time, after which it expands quite aggressively?

These questions are, as of now, still unresolved.

Part IV.

Final Conclusions

Chapter 16

Conclusions and Future Work

To sum up this work, these are the “milestone results” that should be clear:


- In general, physically interesting QFTs have multiple solutions, which are associated and determined by the contours of integration that render the partition function finite, which is absolutely analogous to finding all solutions to the Schwinger-Dyson equations of the problem (and its boundary conditions).
- In order to be able to find all possible solutions, we are forced to extend the field configurations to complex-valued fields, or, more generally, to matrix-valued or Lie-algebra-valued fields. In turn, this implies that the parameters of the theory (mass, coupling constants, etc) are also extended in an analogous fashion.
- The Lee-Yang zeros of these multiple solutions condense over Stokes’ Lines, coalescing the solutions and creating “sectors of analyticity” for the partition function. Thus, we end up having a “canonical” set of superselection rules in order to determine the particular vacuum under study.
- In this sense, the partition function should be understood as a meromorphic function of the parameters of the theory. This is in complete accord with the interpretation of the partition function as a particular D -module, constructed from the 0-dimensional version of the model at hand, in order to engineer a d -dimensional theory. Moreover, once the partition function is being understood as a function of the parameters of theory, which have been “extended”, we can focus on its transformation properties, in particular, whether or not it possess modular symmetry ($SL(2, \mathbb{C})/\mathbb{Z}_2$), which is the automorphism group of the complexified parameter space (understood as a Riemann sphere).
- On a slightly different token, when gauge symmetry is present, the boundary conditions of the Schwinger-Dyson equations (resp. contours that render the partition function finite) and the gauge symmetry completely determine the local system (in the sense of [30]) in question and, in this fashion, they determine some properties of the Higgs bundle associated with this theory. In particular, it is possible to infer the topology of this Higgs bundles, based on the Jacobi metric

of the problem. This information is important for the Geometric Langlands Duality, where the singularity structure of the gauge connection is of importance.

- If we are given a pure gauge theory, $S = \frac{1}{g} \text{tr}(\mathbf{F}^2)$, the contours of the partition function ultimately determine the coupling constant, which makes its appearance in the structure constants determining the Lie algebra of the gauge group at hand (analogously, mathematicians understand this as determining the scalar product of the Lie algebra), thus measuring the “intensity” of the field interactions. In this sense, the structure constants can be dynamically determined by the theory, and its different equivalence classes will determine distinct theories.

Furthermore, in the sense of [57, 63], we can understand the process of starting from a “bare” (initial) theory and dynamically determine its parameters (via the contours of the partition function or the boundary condition of the Schwinger-Dyson equations) giving rise to multiple (final) theories as a *quantization* scheme. This is intimately related to the fact that the quantum corrections, being handles attached to the classical solution, deform the classical geometry into a quantum one — a fact which is related to the more common Geometric and Deformation Quantization schemes.

In terms of cobordisms, this can be picture as shown below, where arrows (which represent the particular propagator in question) going from the initial theory to each one of the final theories represents the quantization of that particular solution:



As for future work, some options are clear, e.g., a better understanding of how the boundary conditions of the Schwinger-Dyson equations determine the structure constants in a pure gauge theory, and how this affects the different multiple solutions of the theory; a deeper realization of which role the Geometric Langlands Duality plays

in this framework; and, of course, a more robust comprehension of the 3-dimensional gravity problem. On the mollifier front, we hope to be able to tackle more physically interesting problems as soon as we have better hardware and software available.

Part V.

Appendices

Convolutions and Smoothing

With the aid of the tools developed below, the smooth approximations of functions can be done in a mathematically rigorous fashion. This is useful to justify the statements made in this work, where this technology is applied to the generating functional of an arbitrary QFT. For proofs of the theorems shown below, see [20] [21].

A.1. Mollifiers

Notation. If $U \subset \mathbb{R}^n$ is open, ∂U is its boundary and $\varepsilon > 0$, let $U_\varepsilon = \{x \in U \mid \text{dist}(x, \partial U) > \varepsilon\}$. Further, let $B(0, \varepsilon)$ be the ball centered on 0 and with radius ε .

Definition. (Standard mollifier.)

A mollifier, η , also called an approximate identity, is a positive $\mathcal{C}^\infty(\mathbb{R}^n)$ function. The standard mollifier is defined in the following way:

- Define $\eta \in \mathcal{C}^\infty(\mathbb{R}^n)$ to be,

$$\eta(x) = \begin{cases} C \exp\left(\frac{1}{|x|^2-1}\right), & \text{if } |x| < 1; \\ 0, & \text{if } |x| \geq 1. \end{cases}$$

The constant $C > 0$ selected so that $\int_{\mathbb{R}^n} \eta(x) dx = 1$.

- $\forall \varepsilon > 0$, set

$$\eta_\varepsilon(x) = \frac{1}{\varepsilon^n} \eta(x/\varepsilon) .$$

η is called the standard mollifier. The functions $\eta_\varepsilon \in \mathcal{C}^\infty$ satisfy $\int_{\mathbb{R}^n} \eta_\varepsilon(x) dx = 1$ and $\text{supp}(\eta_\varepsilon) \subset B(0, \varepsilon)$.

Definition. (Mollification.) If $f : U \rightarrow \mathbb{R}$ is locally integrable, define its mollification to be,

$$U_\varepsilon \ni f^\varepsilon = \eta_\varepsilon * f .$$

That is, $\forall x \in U_\varepsilon$,

$$f^\varepsilon(x) = \int_U \eta_\varepsilon(x-y) f(y) dy = \int_{B(0,\varepsilon)} \eta_\varepsilon(y) f(x-y) dy . \quad (\text{A.1})$$

A.2. Properties of Mollifiers

Theorem A.2.1. (*Properties of mollifiers.*)

1. $f^\varepsilon \in \mathcal{C}^\infty(U_\varepsilon)$;
2. $f^\varepsilon \rightarrow f$, almost everywhere, as $\varepsilon \rightarrow 0$;
3. If $f \in \mathcal{C}(U)$, then $f^\varepsilon \rightarrow f$ uniformly on compact subsets of U ; $\mathcal{E}$
4. If $1 \leq p < \infty$ and $f \in L^p_{\text{loc}}(U)$, then $f^\varepsilon \rightarrow f$ in $L^p_{\text{loc}}(U)$.

Appendix B

Entropy Calculations

In the cases considered in 7.1, the information theoretic entropy can be computed analytically. There are 2 types of functions to be considered:

$$f_{\epsilon}^1(x) = \exp\left\{-\frac{1}{2}\left(\frac{x+c}{\epsilon}\right)^2\right\};$$

$$f_{\epsilon}^2(x) = \exp\left\{-\frac{1}{2}\left(\frac{x^2+c}{\epsilon^2}\right)\right\};$$

where c is a constant scalar. The first type above deals with the Airy Function action for $J < 0$ ($c = \pm\sqrt{-J}$) and the ϕ^4 action with $\mu < 0$ ($c = \pm\sqrt{-\mu/2g}$), while the second type above handles the Airy Function action for $J \geq 0$ ($c = \pm J$) and the ϕ^4 action for $\mu \geq 0$ ($c = \pm i\sqrt{\mu/2g}$).

The information theoretic entropy has the general form,



$$S^i(\epsilon) = -\frac{\int f_{\epsilon}^i(x) \log(f_{\epsilon}^i(x)) dx}{\int f_{\epsilon}^i(x) dx + \log\left(\int f_{\epsilon}^i(x) dx\right)}.$$

Therefore, we see that the analytical results are given by:

$$S^1(\epsilon) = \frac{(\epsilon/2) \sqrt{2\pi}}{\sqrt{2\pi} \epsilon + \log(\sqrt{2\pi} \epsilon)};$$

$$S^2(\epsilon, c) = \frac{e^{-c/2\epsilon^2} \sqrt{2\pi} (\epsilon + c/2\epsilon)}{e^{-c/2\epsilon^2} \sqrt{2\pi} \epsilon + \log(e^{-c/2\epsilon^2} \sqrt{2\pi} \epsilon)}.$$

And, thus, we have closed forms for the entropies of the examples given. The graphs below show these entropies: the leftmost one shows $\mathcal{S}^1(\epsilon)$ while the rightmost one shows $\mathcal{S}^2(\epsilon, c)$ for a range of c values.



Parabolic Cylinder Functions

The differential equation for the Parabolic Cylinder functions is usually written as,

$$\frac{d^2 f}{dz^2} + (a z^2 + b z + c) f = 0 ;$$

where $z \in \mathbb{C}$.

The 0-dimensional ϕ^4 action is given by $S[\phi] = \mu \phi^2/2 + g \phi^4/4$, which yields two Schwinger-Dyson equations:

$$\begin{aligned} \text{Perturbative solution:} & \quad -i \delta_J \mathcal{Z} - J \mathcal{Z} = 0 ; \\ \text{Non-perturbative solutions:} & \quad \delta_J^2 \mathcal{Z} + J \mathcal{Z} - \beta \mathcal{Z} = 0 ; \end{aligned}$$

where $\beta = \mu/g$. The match to the previous form is obtained when $a = 0$, $b = +1$, $c = -\beta$. Note that there are two non-perturbative solutions, one being solitonic and the other one being called simply broken-symmetric.

Part VI.

Bibliography

Bibliography

- [1] Elliott H. Lieb, Michael Loss, *Analysis*, GSM, American Mathematical Society, 2001.
- [2] S. B. Fahy, D. R. Hamann, *Positive-projection Monte Carlo simulation: A new variational approach to strongly interacting fermion systems*, *Phys. Rev. Lett.* **65** (1990) 3437.
- [3] S. B. Fahy, D. R. Hamann, *Diffusive behavior of states in the Hubbard-Stratonovitch transformation*, *Phys. Rev. B* **43** (1991) 765.
- [4] M. Suzuki, *Fractal decomposition of exponential operators with applications to many-body theories and Monte Carlo simulations*, *Phys. Lett. A* **146** (1990) 319.
- [5] P. de Vries, Ph.D. Thesis, University of Amsterdam, 1991, unpublished.
- [6] A. Muramatsu, G. Zumbach, X. Zotos, *A Geometrical View of the Minus-Sign Problem*, *Int. J. Mod. Phys. C* **3** (1992) 185.
- [7] A.H. Zemanian, *Distribution Theory and Transform Analysis: An Introduction to Generalized Functions, with Applications*, Dover, 1987.
- [8] D. Sabo, J.D. Doll, D.L. Freeman, *Stationary Tempering and the Complex Quadrature Problem*, *J. Chem. Phys.* **116** (2002) 3509.
- [9] S. Garcia, Z. Guralnik, G.S. Guralnik, *Theta Vacua and Boundary Conditions of the Schwinger-Dyson Equations*, [[hep-th/9612079](#)]. G.S. Guralnik, Z. Guralnik, *Complexified Path Integrals and the Phases of Quantum Field Theory*, [[arXiv:0710.1256](#)].
- [10] M.G.G. Laidlaw, C.M. DeWitt, *Feynman Functional Integrals for Systems of Indistinguishable Particles*, *Phys. Rev. D* **3** (1971) 1375.
- [11] J.D. Doll, *Notes on the Stationary Phase Monte Carlo Method*, unpublished, 1999.
- [12] R. Easter, D. D. Ferrante, G.S. Guralnik and D. Petrov, *A review of two novel numerical methods in QFT*, [[hep-lat/0306038](#)].

- [13] D. D. Ferrante, J. Doll, G.S. Guralnik, D. Sabo, *Mollified Monte Carlo*, *Nucl. Phys.* **119** (*Proc. Suppl.*) (2003) 965, [[hep-lat/0209053](#)].
- [14] Giovanni Ossola, Alan D. Sokal, *Systematic errors due to linear congruential random-number generators with the Swendsen-Wang algorithm: A warning*, *Phys. Rev.* **E 70** (2004) 027701, [[hep-lat/0403010](#)].
- [15] M. Matsumoto, T. Nishimura, *Mersenne Twister: A 623-dimensionally equidistributed uniform pseudorandom number generator*, *ACM Trans. on Modeling and Computer Simulation* **8** (1998) 3.
- [16] Tuomas J. Lukka, Janne V. Kujala, *Using Genetic Operators to Speed up Markov Chain Monte Carlo Integration*, *Monte Carlo Methods and Appl.* **8** (2002) 51, [[hep-lat/0403010](#)].
- [17] J.C. Baez, *Diffeomorphism invariant generalized measures on the space of connections modulo gauge transformations*, [[hep-th/9305045](#)].
- [18] A. Ashtekar, J. Lewandowski, *Differential geometry on the space of connections via graphs and projective limits*, *J. Geom. Phys.* **17** (1995) 191, [[hep-th/9412073](#)].
- [19] L.D. Landau, E.M. Lifshitz, *Statistical Physics*, Part 1, Butterworth-Heinemann, 1984.
- [20] Lawrence C. Evans, *Partial Differential Equations*, GSM, American Mathematical Society, 1998.
- [21] Michael Reed, Barry Simon, *Methods of Modern Mathematical Physics, Vol II: Fourier Analysis, Self-Adjointness*, Academic Press, 1975.
- [22] Theodore Frankel, *The Geometry of Physics: An Introduction*, Cambridge University Press, 2003.
- [23] M. Nakahara, *Geometry, Topology and Physics*, Taylor & Francis, 2003.
- [24] R. W. R. Darling, *Differential Forms and Connections*, Cambridge University Press, 1994.
- [25] Y. Choquet-Bruhat, C. DeWitt-Morette, *Analysis, Manifolds and Physics, Part I and II*, North Holland.
- [26] Yvan Kerbrat, Helene Kerbrat-Lunc, *Spontaneous symmetry breaking and principal fibre bundles*, *J. Geometry and Physics* Vol. III, No. 2 (1986).
- [27] R. O. Fulp, L. K. Norris, *Splitting of the connection in gauge theories with broken symmetry*, *J. of Mathematical Physics*, Vol. 24, 1871, (1983).
- [28] M. E. Mayer, *The geometry of symmetry breaking in gauge theories*, *Act. Phys. Austr. (Suppl.)* XXIII, 477–490 (1941).

- [29] Gennadi Sardanashvily, *On the geometry of spontaneous symmetry breaking*, *J. Math. Phys.* **33** (4), April 1 1992.
- [30] Carlos T. Simpson, *Higgs bundles and local systems*, *Publications Mathématiques de l'IHÉS*, **75** (1992), p. 5–95.
- [31] Y. Ne'eman, *Geometrization of Spontaneously Broken Gauge Symmetries*, *Theoretical and Mathematical Physics*, **139**(3): 745–750 (2004).
- [32] G. S. Guralnik, C. R. Hagen, T. W. B. Kibble, *Global Conservation Laws and Massless Particles*, *Phys. Rev. Lett.* **13**, 585–587 (1964). P. W. Higgs, *Broken symmetries, massless particles and gauge fields*, *Phys. Lett.* **12** 132. F. Englert and R. Brout, *Broken Symmetry and the Mass of Gauge Vector Mesons*, *Phys. Rev. Lett.* **13**, 321–323 (1964). P. W. Higgs, *Broken Symmetries and the Masses of Gauge Bosons*, *Phys. Rev. Lett.* **13**, 508–509 (1964).
- [33] F. Strocchi, *Spontaneous Symmetry Breaking in Local Gauge Quantum Field Theory; The Higgs Mechanism*, *Commun. Math. Phys.* **56**, 57–78 (1977).
- [34] Sidney Coleman, Erick Weinberg, *Radiative Corrections as the Origin of Spontaneous Symmetry Breaking*, *Phys. Rev. D* **7**, 1888–1910 (1973). Erick Weinberg, *Radiative Corrections as the Origin of Spontaneous Symmetry Breaking*, [hep-th/0507214](https://arxiv.org/abs/hep-th/0507214).
- [35] S. Garcia, G. Guralnik, Z. Guralnik, *Theta Vacua and Boundary Conditions of the Schwinger Dyson Equations*, [hep-th/9612079](https://arxiv.org/abs/hep-th/9612079). G. S. Guralnik, Z. Guralnik, *Complexified Path Integrals and the Phases of Quantum Field Theory*, [[arXiv:0710.1256](https://arxiv.org/abs/0710.1256)].
- [36] R. E. Borcherds and A. Barnard, *Lectures on Quantum Field Theory*, [math-ph/0204014](https://arxiv.org/abs/math-ph/0204014).
- [37] M. V. Berry, *Stokes' phenomenon; smoothing a victorian discontinuity*, *Publications Mathématiques de l'IHÉS*, **68** (1988), p. 211–221. M. V. Berry, *Uniform Asymptotic Smoothing of Stokes's Discontinuities*, *Proc. R. Soc. Lond. A*, Vol. **422**, No. 1862 (Mar. 8, 1989), pp. 7–21.
- [38] J. Milne, *Modular Functions and Modular Forms*, <http://www.jmilne.org/math/CourseNotes/math678.html>
- [39] E. Witten, *Three-Dimensional Gravity Revisited*, [[arXiv:0706.3359](https://arxiv.org/abs/0706.3359)]. A. Maloney, E. Witten, *Quantum Gravity Partition Functions in Three Dimensions*, [[arXiv:0712.0155](https://arxiv.org/abs/0712.0155)].
- [40] D. D. Ferrante, G. S. Guralnik, *Mollifying Quantum Field Theory or Lattice QFT in Minkowski Spacetime and Symmetry Breaking*, [[arXiv:hep-lat/0602013](https://arxiv.org/abs/hep-lat/0602013)].
- [41] D. D. Ferrante, G. S. Guralnik, *From Symmetry Breaking to Topology Change I*, [[arXiv:hep-th/0609190](https://arxiv.org/abs/hep-th/0609190)].

- [42] Klaus Fredenhagen and Karl-Henning Rehren and Erhard Seiler, *Quantum Field Theory: Where We Are*, [hep-th/0603155](#).
- [43] M. Reed and B. Simon; *Methods of modern mathematical physics*, Volume 1: Functional Analysis. Academic Press, New York 1972.
- [44] N. Dunford and J. T. Schwartz ; *Linear operators*, Volume 2: Spectral Theory. Interscience, New York 1964.
- [45] Sidney Coleman, Frank De Luccia, *Gravitational effects on and of vacuum decay*, [Phys. Rev. D 21, 3305–3315 \(1980\)](#).
- [46] Detlef Laugwitz, *Differential and Riemannian Geometry*, Academic Press, 1965. Cornelius Lanczos, *The Variational Principles of Mechanics*, Dover Publications, 1986. V. I. Arnold, *Mathematical Methods of Classical Mechanics*. Springer, 1997.
- [47] N. Seiberg, E. Witten, *Monopoles, Duality and Chiral Symmetry Breaking in $N = 2$ Supersymmetric QCD*, Nucl. Phys. B 431 (1994) 484–550, [hep-th/9408099](#).
- [48] S. K. Donaldson, *The Seiberg-Witten equations and 4-manifold topology*, [Bull. Amer. Math. Soc. 33 \(1996\), 45–70](#).
- [49] Liviu I. Nicolaescu, *Notes on Seiberg-Witten Theory*. American Mathematical Society, 2000.
- [50] Jürgen Jost, *Riemannian Geometry and Geometric Analysis*. Springer.
- [51] Marek Szydłowski, Michale Heller, Wiesław Sasin, *Geometry of spaces with the Jacobi metric*, [J. Math. Phys. 37 \(1\), 1996, 346–360](#).
- [52] E. Witten, *Mirror Symmetry, Hitchin's Equations, And Langlands Duality*, [arXiv:0802.0999](#).
- [53] M. Kontsevich, *Intersection theory on the moduli space of curves and the matrix Airy function*, [Comm. Math. Phys. Volume 147, Number 1 \(1992\), 1-23](#). R. N. Fernandez, V. S. Varadarajan, *Airy Functions for Compact Lie Groups*, [[arXiv:0707.3235](#)].
- [54] Harish-Chandra, *Differential operators on a semisimple Lie algebra*, [Am. J. Math. 79, \(1957\) 87-120](#). Harish-Chandra, *Some applications of invariant differential operators on a semi simple Lie algebra*, [Séminaire Bourbaki, 4 \(1956-1958\), Exposé No. 160, 8 p.](#)
- [55] S. Gukov, E. Witten, *Gauge Theory, Ramification, And The Geometric Langlands Program*, [[arXiv:hep-th/0612073](#)]; E. Witten, *Gauge Theory And Wild Ramification*, [[arXiv:0710.0631](#)].
- [56] C. Vafa, *Geometric Physics*, [[arXiv:hep-th/9810149](#)].

- [57] T. Eguchi, P. Gilkey, A. J. Hanson, *Gravitation, gauge theories and differential geometry*, *Phys. Rept.* **66** (1980) 213.
- [58] M. Atiyah, *Geometry of Yang-Mills Fields*, *Lecture Notes in Physics* **80** (1978) 216.
- [59] K. Hori, S. Katz, A. Klemm, R. Pandharipande, R. Thomas, C. Vafa, R. Vakil, E. Zaslow, *Mirror Symmetry*, *Clay Mathematics Monographs* (2003).
- [60] A. Alexandrov, A. Mironov, A. Morozov, *Partition Functions of Matrix Models as the First Special Functions of String Theory I. Finite Size Hermitean 1-Matrix Model*, [[arXiv:hep-th/0310113](#)]. L. Chekhov, A. Marshakov, A. Mironov, D. Vasiliev, *Complex Geometry of Matrix Models*, [[arXiv:hep-th/0506075](#)]. A. Mironov, *Matrix Models vs. Matrix Integrals*, [[arXiv:hep-th/0506158](#)]. A. Alexandrov, A. Mironov, A. Morozov, *M-Theory of Matrix Models*, [[arXiv:hep-th/0605171](#)]. A. Mironov, A. Morozov, *Virasoro constraints for Kontsevich-Hurwitz partition function*, [[arXiv:0807.2843](#)]. A. Morozov, Sh. Shakirov, *Analogue of the identity $\text{Log Det} = \text{Trace Log}$ for resultants*, [[arXiv:0804.4632](#)]. A. Morozov, Sh. Shakirov, *Resultants and Contour Integrals*, [[arXiv:0807.4539](#)].
- [61] L. Saïdane, *Classe d'équivalence Formelle D'un D-Module D'Airy*, [[arXiv:math/0306170](#)].
- [62] S. Sternberg, *Toronto Lectures on Physics*.
- [63] M. F. Atiyah, *Topological quantum field theory*, *Publications Mathématiques de l'IHÉS*, **68** (1988), p. 175-186. G. B. Segal, *The definition of conformal field theory*, *Proc. 16th Int. Conf., NATO Adv. Res. Workshop, Como/Italy 1987, NATO ASI Ser., Ser. C 250*, 165-171 (1988). U. Schreiber, *AQFT from n-functorial QFT*, [[arXiv:0806.1079](#)].